

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
12 June 2003 (12.06.2003)

PCT

(10) International Publication Number  
WO 03/047595 A1

- (51) International Patent Classification<sup>7</sup>: A61K 31/59 (74) Agents: ESMOND, Robert, W. et al.; Sterne, Kessler, Goldstein & Fox P.L.L.C., 1100 New York Avenue, N.W., Suite 600, Washington, DC 20005-3934 (US).
- (21) International Application Number: PCT/US02/38483
- (22) International Filing Date: 3 December 2002 (03.12.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/334,554 3 December 2001 (03.12.2001) US
- (71) Applicant: NOVACEA, INC. [US/US]; 601 Gateway Boulevard, Suite 450, South San Francisco, CA 94080 (US).
- (71) Applicants and  
(72) Inventors: CHEN, Andrew-Xian [US/US]; 4646 Bryson Terrace, San Diego, CA 92130 (US). FAN, Jun [CN/US]; 10566 Giffin Way, San Diego, CA 92126 (US). YU, Xi-Yun [CN/US]; 11441 Creekstone Lane, San Diego, CA 92128 (US). WHITEHOUSE, Martha, J. [US/US]; 151 Upper Terrace, San Francisco, CA 94117 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PHARMACEUTICAL COMPOSITIONS COMPRISING ACTIVE VITAMIN D COMPOUNDS

(57) Abstract: Disclosed are pharmaceutical compositions comprising an active vitamin D compound in emulsion pre-concentrate formulations, as well as emulsions and sub-micron droplet emulsions produced therefrom. The compositions comprise a lipophilic phase component, one or more surfactants, and an active vitamin D compound. The compositions may optionally further comprise a hydrophilic phase component.

WO 03/047595 A1

PHARMACEUTICAL COMPOSITIONS COMPRISING  
ACTIVE VITAMIN D COMPOUNDS

BACKGROUND OF THE INVENTION

Field of the Invention

[0001] The present invention relates to novel pharmaceutical compositions comprising an active vitamin D compound, wherein the pharmaceutical compositions are emulsion pre-concentrates. The invention also relates to emulsions and sub-micron droplet emulsions produced upon dilution of the emulsion pre-concentrates with an aqueous solution.

Background Art

[0002] Vitamin D is a fat soluble vitamin which is essential as a positive regulator of calcium homeostasis. (See Harrison's Principles of Internal Medicine: Part Eleven, "Disorders of Bone and Mineral Metabolism," Chapter 335, pp. 1860-1865, E. Braunwald *et al.*, (eds.), McGraw-Hill, New York (1987)). The active form of vitamin D is  $1\alpha,25$ -dihydroxyvitamin  $D_3$ , also known as calcitriol. Specific nuclear receptors for active vitamin D compounds have been discovered in cells from diverse organs not involved in calcium homeostasis. (Miller *et al.*, *Cancer Res.* 52:515-520 (1992)). In addition to influencing calcium homeostasis, active vitamin D compounds have been implicated in osteogenesis, modulation of immune response, modulation of the process of insulin secretion by the pancreatic B cell, muscle cell function, and the differentiation and growth of epidermal and hematopoietic tissues.

[0003] Moreover, there have been many reports demonstrating the utility of active vitamin D compounds in the treatment of cancer. For example, it has been shown that certain vitamin D compounds and analogues possess potent antileukemic activity by virtue of inducing the differentiation of malignant cells (specifically, leukemic cells) to non-malignant macrophages (monocytes) and are useful in the treatment of leukemia. (Suda *et al.*, U.S. Patent No. 4,391,802; Partridge *et al.*, U.S. Patent No. 4,594,340). Antiproliferative and differentiating

actions of calcitriol and other vitamin D<sub>3</sub> analogues have also been reported with respect to the treatment of prostate cancer. (Bishop *et al.*, U.S. Patent No. 5,795,882). Active vitamin D compounds have also been implicated in the treatment of skin cancer (Chida *et al.*, *Cancer Research* 45:5426-5430 (1985)), colon cancer (Disman *et al.*, *Cancer Research* 47:21-25 (1987)), and lung cancer (Sato *et al.*, *Tohoku J. Exp. Med.* 138:445-446 (1982)). Other reports suggesting important therapeutic uses of active vitamin D compounds are summarized in Rodriguez *et al.*, U.S. Patent No. 6,034,079.

[0004] Although the administration of active vitamin D compounds may result in substantial therapeutic benefits, the treatment of cancer and other diseases with such compounds is limited by the effects these compounds have on calcium metabolism. At the levels required *in vivo* for effective use as anti-proliferative agents, active vitamin D compounds can induce markedly elevated and potentially dangerous blood calcium levels by virtue of their inherent calcemic activity. That is, the clinical use of calcitriol and other active vitamin D compounds as anti-proliferative agents is precluded, or severely limited, by the risk of hypercalcemia.

[0005] It has been shown that the problem of systemic hypercalcemia can be overcome by "pulse-dose" administration of a sufficient dose of an active vitamin D compound such that an anti-proliferative effect is observed while avoiding the development of severe hypercalcemia. (WO 99/49870). According to WO 99/49870, the active vitamin D compound may be administered no more than every three days, for example, once a week at a dose of at least 0.12 µg/kg per day (8.4 µg in a 70 kg person). Pharmaceutical compositions used in the pulse-dose regimen of WO 99/49870 comprise 5-100 µg of active vitamin D compound and may be administered in the form for oral, intravenous, intramuscular, topical, transdermal, sublingual, intranasal, intratumoral or other preparations.

[0006] ROCALTROL is the trade name of a calcitriol formulation sold by Roche Laboratories. ROCALTROL is available in the form of capsules containing 0.25 and 0.5 µg calcitriol and as an oral solution containing 1 µg/mL of calcitriol. All dosage forms contain butylated hydroxyanisole (BHA) and butylated

hydroxytoluene (BHT) as antioxidants. The capsules also contain a fractionated triglyceride of coconut oil and the oral solution contains a fractionated triglyceride of palm seed oil. (Physician's Desk Reference, 54<sup>th</sup> Edition, pp 2649-2651, Medical Economics Company, Inc., Montvale, NJ (2000)).

[0007] It is known that calcitriol is light-sensitive and is especially prone to oxidation. Moreover, calcitriol and other active vitamin D compounds are lipophilic, meaning that they are soluble in lipids and some organic solvents, while being substantially insoluble or only sparsely soluble in water. Because of the lipophilic nature of active vitamin D compounds, the dispersion of such compounds in aqueous solutions, such as the gastric fluids of the stomach, is significantly limited. Accordingly, the pharmacokinetic parameters of active vitamin D compound formulations heretofore described in the art are sub-optimal for use with high dose pulse administration regimens. In addition, the active vitamin D compound formulations that are currently available tend to exhibit substantial variability of absorption in the small intestine. Moreover, for oral administration, the relationship between dosage and blood concentration that is observed with most active vitamin D compound formulations is not linear; that is, the quantity of compound absorbed into the blood stream does not correlate with the amount of compound that is administered in a given dose, especially at higher dosage levels.

[0008] Thus, there is a need for improved pharmaceutical compositions comprising active vitamin D compounds, particularly in the context of pulse-dose treatment regimens that are designed to provide anti-proliferative (*e.g.*, anti-cancer) benefits while avoiding the consequence of hypercalcemia. In particular, a need exists in the art for a pharmaceutical composition comprising an active vitamin D compound that remains stable over prolonged periods of time, even at elevated temperatures, while at the same time exhibiting improved pharmacokinetic parameters for the active vitamin D compound, and reduced variability in absorption, when administered to a patient.

## BRIEF SUMMARY OF THE INVENTION

[0009] The present invention overcomes the disadvantages heretofore encountered in the art by providing pharmaceutical compositions comprising active vitamin D compounds in emulsion pre-concentrate formulations. The pharmaceutical compositions of the present invention are an advance over the prior art in that they provide a dosage form of active vitamin D compounds, such as calcitriol, in a sufficiently high concentration to permit convenient use, stability and rapid dispersion in solution, and yet meet the required criteria in terms of pharmacokinetic parameters, especially in the context of pulse-dosing administration regimens. More specifically, in a preferred embodiment, the pharmaceutical compositions of the present invention exhibit a  $C_{\max}$  that is at least 1.5 to two times greater than the  $C_{\max}$  that is observed with ROCALTROL, and a shorter  $T_{\max}$  than that which is observed with ROCALTROL.

[0010] The emulsion pre-concentrates of the present invention are non-aqueous formulations for an active vitamin D compound that are capable of providing a pharmaceutically acceptable emulsion, upon contact with water or other aqueous solution.

[0011] According to one aspect of the invention, pharmaceutical compositions are provided comprising (a) a lipophilic phase component, (b) one or more surfactants, and (c) an active vitamin D compound; wherein said composition is an emulsion pre-concentrate, which upon dilution with water in a water to composition ratio of about 1:1 or more of water forms an emulsion having an absorbance of greater than 0.3 at 400 nm. According to this aspect of the invention, the pharmaceutical compositions may further comprise a hydrophilic phase component.

[0012] According to another aspect of the invention, a pharmaceutical emulsion composition is provided comprising water and an emulsion pre-concentrate, said emulsion pre-concentrate comprising (a) a lipophilic phase component, (b) one or more surfactants, and (c) an active vitamin D compound, and optionally, a hydrophobic phase component.

[0013] The emulsions produced from the emulsion pre-concentrates of the present invention (upon dilution with water) include both emulsions as conventionally understood by those of ordinary skill in the art (*i.e.*, a dispersion of an organic phase in water); as well as "sub-micron droplet emulsions" (*i.e.*, dispersions of an organic phase in water wherein the average diameter of the dispersion particles is less than 1000 nm.)

[0014] According to another aspect of the invention, methods are provided for the preparation of emulsion pre-concentrates comprising active vitamin D compounds. The methods encompassed within this aspect of the invention comprise bringing an active vitamin D compound, *e.g.*, calcitriol, into intimate admixture with a lipophilic phase component and with one or more surfactants, and optionally, with a hydrophilic phase component.

[0015] In yet another aspect of the invention, methods are provided for the treatment and prevention of hyperproliferative diseases such as cancer and psoriasis, said methods comprising administering an active vitamin D compound in an emulsion pre-concentrate formulation to a patient in need thereof. Alternatively, the active vitamin D compound can be administered in an emulsion formulation that is made by diluting an emulsion pre-concentrate of the present invention with an appropriate quantity of water. In a preferred embodiment of this aspect of the invention, the administration of the active vitamin D compound to a patient is accomplished by using, *e.g.*, a pulse dosing regimen. For example, according to this aspect of the invention, an active vitamin D compound in an emulsion pre-concentrate formulation is administered to a patient no more than once every three days at a dose of at least 0.12  $\mu\text{g/kg}$  per day.

## BRIEF DESCRIPTIONS OF THE FIGURES

- [0016] Fig. 1 is a graphical representation of the mean plasma concentration of calcitriol in dogs versus time following administration of three different formulations of calcitriol at a dose of 1 µg/kg.
- [0017] Figs. 2A and 2B are graphical representations of the mean plasma concentration-time curve for calcitriol after escalating doses of semi-solid #3 in male (Fig. 2A) and female (Fig. 2B) dogs.
- [0018] Figs. 3A and 3B are graphical representations of the plasma concentration-time curve for calcitriol in male (Fig. 3A) and female (Fig. 3B) dogs after semi-solid #3 dosing.
- [0019] Figs. 4A and 4B are graphical representations of the mean serum calcium after increasing doses of semi-solid #3 in male (Fig. 4A) and female (Fig. 4B) dogs.
- [0020] Figs. 5A-5C are graphical representations of the plasma calcitriol and serum calcium data following administration of semi-solid #3 in male dogs.
- [0021] Fig. 6 is a graphical representation of the mean plasma concentration of calcitriol by dose group in humans following administration of semi-solid #3.

## DETAILED DESCRIPTION OF THE INVENTION

- [0022] The present invention is directed to pharmaceutical compositions comprising active vitamin D compounds in emulsion pre-concentrate formulations. The compositions of the invention meet or substantially reduce the difficulties associated with active vitamin D compound therapy hitherto encountered in the art including, in particular, undesirable pharmacokinetic parameters of the compound upon administration to a patient.
- [0023] It has been found that the compositions of the invention permit the preparation of semi-solid and liquid compositions containing an active vitamin D compound in sufficiently high concentration to permit, *e.g.*, convenient oral

administration, while at the same time achieving improved pharmacokinetic parameters for the active vitamin D compound. For example, as compared to ROCALTROL, the compositions of the present invention exhibit a  $C_{\max}$  that is at least 1.5 to two times greater than the  $C_{\max}$  that is observed with ROCALTROL, and a shorter  $T_{\max}$  than that which is observed with ROCALTROL. Preferably, the pharmaceutical compositions of the present invention provide a  $C_{\max}$  of at least about 900 pg/mL plasma, more preferably about 900 to about 3000 pg/mL plasma, more preferably about 1500 to about 3000 pg/mL plasma. In addition, the compositions of the invention preferably provide a  $T_{\max}$  of less than about 6.0 hours, more preferably about 1.0 to about 3.0 hours, more preferably about 1.5 to about 2.0 hours. In addition, the compositions of the invention preferably provide a  $T_{1/2}$  of less than about 25 hours, more preferably about 2 to about 10 hours, more preferably about 5 to about 9 hours.

[0024] The term  $C_{\max}$  is defined as the maximum concentration of active vitamin D compound achieved in the serum following administration of the drug. The term  $T_{\max}$  is defined as the time at which  $C_{\max}$  is achieved. The term  $T_{1/2}$  is defined as the time required for the concentration of active vitamin D compound in the serum to decrease by half. The disclosed values for pharmacokinetic data apply to the population of recipients of a composition comprising an active vitamin D compound as a whole, not individual recipients. Thus, any individual receiving a composition of the present invention may not necessarily achieve the preferred pharmacokinetic parameters. However, when a composition of the present invention is administered to a sufficiently large population of subjects, the pharmacokinetic parameters will approximately match the values disclosed herein.

[0025] According to one aspect of the present invention, a pharmaceutical composition is provided comprising (a) a lipophilic phase component, (b) one or more surfactants, (c) an active vitamin D compound; wherein said composition is an emulsion pre-concentrate, which upon dilution with water, in a water to composition ratio of about 1:1 or more of said water, forms an emulsion having



an absorbance of greater than 0.3 at 400 nm. The pharmaceutical composition of the invention may further comprise a hydrophilic phase component.

[0026] In another aspect of the invention, a pharmaceutical emulsion composition is provided comprising water (or other aqueous solution) and an emulsion pre-concentrate.

[0027] The term "emulsion pre-concentrate," as used herein, is intended to mean a system capable of providing an emulsion upon contacting with, *e.g.*, water. The term "emulsion," as used herein, is intended to mean a colloidal dispersion comprising water and organic components including hydrophobic (lipophilic) organic components. The term "emulsion" is intended to encompass both conventional emulsions, as understood by those skilled in the art, as well as "sub-micron droplet emulsions," as defined immediately below.

[0028] The term "sub-micron droplet emulsion," as used herein is intended to mean a dispersion comprising water and organic components including hydrophobic (lipophilic) organic components, wherein the droplets or particles formed from the organic components have an average maximum dimension of less than about 1000 nm.

[0029] Sub-micron droplet emulsions are identifiable as possessing one or more of the following characteristics. They are formed spontaneously or substantially spontaneously when their components are brought into contact, that is without substantial energy supply, *e.g.*, in the absence of heating or the use of high shear equipment or other substantial agitation.

[0030] The particles of a sub-micron droplet emulsion may be spherical, though other structures are feasible, *e.g.* liquid crystals with lamellar, hexagonal or isotropic symmetries. Generally, sub-micron droplet emulsions comprise droplets or particles having a maximum dimension (*e.g.*, average diameter) of between about 50 nm to about 1000 nm, and preferably between about 200 nm to about 300 nm.

[0031] The term "pharmaceutical composition" as used herein is to be understood as defining compositions of which the individual components or ingredients are themselves pharmaceutically acceptable, *e.g.*, where oral administration is

foreseen, acceptable for oral use and, where topical administration is foreseen, topically acceptable.

[0032] The pharmaceutical compositions of the present invention will generally form an emulsion upon dilution with water. The emulsion will form according to the present invention upon the dilution of an emulsion pre-concentrate with water in a water to composition ratio of about 1:1 or more of said water. According to the present invention, the ratio of water to composition can be, *e.g.*, between 1:1 and 5000:1. For example, the ratio of water to composition can be about 1:1, 2:1, 3:1, 4:1, 5:1, 10:1, 200:1, 300:1, 500:1, 1000:1, or 5000:1. The skilled artisan will be able to readily ascertain the particular ratio of water to composition that is appropriate for any given situation or circumstance.

[0033] According to the present invention, upon dilution of said emulsion pre-concentrate with water, an emulsion will form having an absorbance of greater than 0.3 at 400 nm. The absorbance at 400 nm of the emulsions formed upon 1:100 dilution of the emulsion pre-concentrates of the present invention can be, *e.g.*, between 0.3 and 4.0. For example, the absorbance at 400 nm can be, *e.g.*, about 0.4, 0.5, 0.6, 1.0, 1.2, 1.6, 2.0, 2.2, 2.4, 2.5, 3.0, or 4.0. Methods for determining the absorbance of a liquid solution are well known by those in the art. The skilled artisan will be able to ascertain and adjust the relative proportions of the ingredients of the emulsions pre-concentrates of the invention in order to obtain, upon dilution with water, an emulsion having any particular absorbance encompassed within the scope of the invention.

[0034] The pharmaceutical compositions of the present invention can be, *e.g.*, in a semi-solid formulation or in a liquid formulation. Semi-solid formulations of the present invention can be any semi-solid formulation known by those of ordinary skill in the art, including, *e.g.*, gels, pastes, creams and ointments.

[0035] The pharmaceutical compositions of the present invention comprise a lipophilic phase component. Suitable components for use as lipophilic phase components include any pharmaceutically acceptable solvent which is non-miscible with water. Such solvents will appropriately be devoid or substantially devoid of surfactant function.

[0036] The lipophilic phase component may comprise mono-, di- or triglycerides.

Mono-, di- and triglycerides that may be used within the scope of the invention include those that are derived from C<sub>6</sub>, C<sub>8</sub>, C<sub>10</sub>, C<sub>12</sub>, C<sub>14</sub>, C<sub>16</sub>, C<sub>18</sub>, C<sub>20</sub> and C<sub>22</sub> fatty acids. Exemplary diglycerides include, in particular, diolein, dipalmitolein, and mixed caprylin-caprin diglycerides. Preferred triglycerides include vegetable oils, fish oils, animal fats, hydrogenated vegetable oils, partially hydrogenated vegetable oils, synthetic triglycerides, modified triglycerides, fractionated triglycerides, medium and long-chain triglycerides, structured triglycerides, and mixtures thereof.

[0037] Among the above-listed triglycerides, preferred triglycerides include: almond oil; babassu oil; borage oil; blackcurrant seed oil; canola oil; castor oil; coconut oil; corn oil; cottonseed oil; evening primrose oil; grapeseed oil; groundnut oil; mustard seed oil; olive oil; palm oil; palm kernel oil; peanut oil; rapeseed oil; safflower oil; sesame oil; shark liver oil; soybean oil; sunflower oil; hydrogenated castor oil; hydrogenated coconut oil; hydrogenated palm oil; hydrogenated soybean oil; hydrogenated vegetable oil; hydrogenated cottonseed and castor oil; partially hydrogenated soybean oil; partially soy and cottonseed oil; glyceryl tricaproate; glyceryl tricaprilate; glyceryl tricaprinate; glyceryl triundecanoate; glyceryl trilaurate; glyceryl trioleate; glyceryl trilinoleate; glyceryl trilinolenate; glyceryl tricaprilate/caprinate; glyceryl tricaprilate/caprinate/laurate; glyceryl tricaprilate/caprinate/linoleate; and glyceryl tricaprilate/caprinate/stearate.

[0038] A preferred triglyceride is the medium chain triglyceride available under the trade name LABRAFAC CC. Other preferred triglycerides include neutral oils, *e.g.*, neutral plant oils, in particular fractionated coconut oils such as known and commercially available under the trade name MIGLYOL, including the products: MIGLYOL 810; MIGLYOL 812; MIGLYOL 818; and CAPTEX 355.

[0039] Also suitable are caprylic-capric acid triglycerides such as known and commercially available under the trade name MYRITOL, including the product MYRITOL 813. Further suitable products of this class are CAPMUL MCT, CAPTEX 200, CAPTEX 300, CAPTEX 800, NEOBEE M5 and MAZOL 1400.

- [0040] Especially preferred as lipophilic phase component is the product MIGLYOL 812. (See U.S. Patent No. 5,342,625).
- [0041] Pharmaceutical compositions of the present invention may further comprise a hydrophilic phase component. The hydrophilic phase component may comprise, *e.g.*, a pharmaceutically acceptable  $C_{1-5}$  alkyl or tetrahydrofurfuryl di- or partial-ether of a low molecular weight mono- or poly-oxy-alkanediol. Suitable hydrophilic phase components include, *e.g.*, di- or partial-, especially partial-, -ethers of mono- or poly-, especially mono- or di-, -oxy-alkanediols comprising from 2 to 12, especially 4 carbon atoms. Preferably the mono- or poly-oxy-alkanediol moiety is straight-chained. Exemplary hydrophilic phase components for use in relation to the present invention are those known and commercially available under the trade names TRANSCUTOL and COLYCOFUROL. (See U.S. Patent No. 5,342,625).
- [0042] In an especially preferred embodiment, the hydrophilic phase component comprises 1,2-propyleneglycol.
- [0043] The hydrophilic phase component of the present invention may of course additionally include one or more additional ingredients. Preferably, however, any additional ingredients will comprise materials in which the active vitamin D compound is sufficiently soluble, such that the efficacy of the hydrophilic phase as an active vitamin D compound carrier medium is not materially impaired. Examples of possible additional hydrophilic phase components include lower (*e.g.*,  $C_{1-3}$ ) alkanols, in particular ethanol.
- [0044] Pharmaceutical compositions of the present invention also comprise one or more surfactants. Surfactants that can be used in conjunction with the present invention include hydrophilic or lipophilic surfactants, or mixtures thereof. Especially preferred are non-ionic hydrophilic and non-ionic lipophilic surfactants.
- [0045] Suitable hydrophilic surfactants include reaction products of natural or hydrogenated vegetable oils and ethylene glycol, *i.e.* polyoxyethylene glycolated natural or hydrogenated vegetable oils, for example polyoxyethylene glycolated natural or hydrogenated castor oils. Such products may be obtained in known

manner, *e.g.*, by reaction of a natural or hydrogenated castor oil or fractions thereof with ethylene oxide, *e.g.*, in a molar ratio of from about 1:35 to about 1:60, with optional removal of free polyethyleneglycol components from the product, *e.g.*, in accordance with the methods disclosed in German Auslegeschriften 1,182,388 and 1,518,819.

[0046] Suitable hydrophilic surfactants for use in the present pharmaceutical compounds also include polyoxyethylene-sorbitan-fatty acid esters, *e.g.*, mono- and triauryl, palmityl, stearyl and oleyl esters, *e.g.*, of the type known and commercially available under the trade name TWEEN; including the products: TWEEN 20 (polyoxyethylene(20)sorbitanmonolaurate), TWEEN 40 (polyoxyethylene(20)sorbitanmonopalmitate), TWEEN 60 (polyoxyethylene(20)sorbitanmonostearate), TWEEN 80 (polyoxyethylene(20)sorbitanmonooleate), TWEEN 65 (polyoxyethylene(20)sorbitantristearate), TWEEN 85 (polyoxyethylene(20)sorbitantrioleate), TWEEN 21 (polyoxyethylene(4)sorbitanmonolaurate), TWEEN 61 (polyoxyethylene(4)sorbitanmonostearate), and TWEEN 81 (polyoxyethylene(5)sorbitanmonooleate).

[0047] Especially preferred products of this class for use in the compositions of the invention are the above products TWEEN 40 and TWEEN 80. (*See* Hauer, *et al.*, U.S. Patent No. 5,342,625).

[0048] Also suitable as hydrophilic surfactants for use in the present pharmaceutical compounds are polyoxyethylene alkylethers; polyoxyethylene glycol fatty acid esters, for example polyoxyethylene stearic acid esters; polyglycerol fatty acid esters; polyoxyethylene glycerides; polyoxyethylene vegetable oils; polyoxyethylene hydrogenated vegetable oils; reaction mixtures of polyols and, *e.g.*, fatty acids, glycerides, vegetable oils, hydrogenated vegetable oils, and sterols; polyoxyethylene-polyoxypropylene co-polymers; polyoxyethylene-polyoxypropylene block co-polymers; dioctylsuccinate, dioctylsodiumsulfosuccinate, di-[2-ethylhexyl]-succinate or sodium lauryl sulfate; phospholipids, in particular lecithins such as, *e.g.*, soya bean lecithins; propylene

glycol mono- and di-fatty acid esters such as, *e.g.*, propylene glycol dicaprylate, propylene glycol dilaurate, propylene glycol hydroxystearate, propylene glycol isostearate, propylene glycol laurate, propylene glycol ricinoleate, propylene glycol stearate, and, especially preferred, propylene glycol caprylic-capric acid diester; and bile salts, *e.g.*, alkali metal salts, for example sodium taurocholate.

[0049] Suitable lipophilic surfactants include alcohols; polyoxyethylene alkylethers; fatty acids; bile acids; glycerol fatty acid esters; acetylated glycerol fatty acid esters; lower alcohol fatty acids esters; polyethylene glycol fatty acids esters; polyethylene glycol glycerol fatty acid esters; polypropylene glycol fatty acid esters; polyoxyethylene glycerides; lactic acid esters of mono/diglycerides; propylene glycol diglycerides; sorbitan fatty acid esters; polyoxyethylene sorbitan fatty acid esters; polyoxyethylene-polyoxypropylene block copolymers; trans-esterified vegetable oils; sterols; sugar esters; sugar ethers; sucroglycerides; polyoxyethylene vegetable oils; polyoxyethylene hydrogenated vegetable oils; reaction mixtures of polyols and at least one member of the group consisting of fatty acids, glycerides, vegetable oils, hydrogenated vegetable oils, and sterols; and mixtures thereof.

[0050] Suitable lipophilic surfactants for use in the present pharmaceutical compounds also include trans-esterification products of natural vegetable oil triglycerides and polyalkylene polyols. Such trans-esterification products are known in the art and may be obtained *e.g.*, in accordance with the general procedures described in U.S. Pat. No. 3,288,824. They include trans-esterification products of various natural (*e.g.*, non-hydrogenated) vegetable oils for example, maize oil, kernel oil, almond oil, ground nut oil, olive oil and palm oil and mixtures thereof with polyethylene glycols, in particular polyethylene glycols having an average molecular weight of from 200 to 800. Preferred are products obtained by trans-esterification of 2 molar parts of a natural vegetable oil triglyceride with one molar part of polyethylene glycol (*e.g.*, having an average molecular weight of from 200 to 800). Various forms of trans-esterification products of the defined class are known and commercially available under the trade name LABRAFIL.

[0051] Additional lipophilic surfactants that are suitable for use with the present pharmaceutical compositions include oil-soluble vitamin derivatives, *e.g.*, tocopherol PEG-1000 succinate ("vitamin E TPGS").

[0052] Also suitable as lipophilic surfactants for use in the present pharmaceutical compounds are mono-, di- and mono/di-glycerides, especially esterification products of caprylic or capric acid with glycerol; sorbitan fatty acid esters; pentaerythritol fatty acid esters and polyalkylene glycol ethers, for example pentaerythrite- -dioleate, -distearate, -monolaurate, -polyglycol ether and -monostearate as well as pentaerythrite-fatty acid esters; monoglycerides, *e.g.*, glycerol monooleate, glycerol monopalmitate and glycerol monostearate; glycerol triacetate or (1,2,3)-triacetin; and sterols and derivatives thereof, for example cholesterol and derivatives thereof, in particular phytosterols, *e.g.*, products comprising sitosterol, campesterol or stigmasterol, and ethylene oxide adducts thereof, for example soya sterols and derivatives thereof.

[0053] It is understood by those of ordinary skill in the art that several commercial surfactant compositions contain small to moderate amounts of triglycerides, typically as a result of incomplete reaction of a triglyceride starting material in, for example, a trans-esterification reaction. Thus, the surfactants that are suitable for use in the present pharmaceutical compositions include those surfactants that contain a triglyceride. Examples of commercial surfactant compositions containing triglycerides include some members of the surfactant families GELUCIRES, MAISINES, AND IMWITORS. Specific examples of these compounds are GELUCIRE 44/14 (saturated polyglycolized glycerides); GELUCIRE 50/13 (saturated polyglycolized glycerides); GELUCIRE 53/10 (saturated polyglycolized glycerides); GELUCIRE 33/01 (semi-synthetic triglycerides of C<sub>8</sub>-C<sub>18</sub> saturated fatty acids); GELUCIRE 39/01 (semi-synthetic glycerides); other GELUCIRE, such as 37/06, 43/01, 35/10, 37/02, 46/07, 48/09, 50/02, 62/05, etc.; MAISINE 35-I (linoleic glycerides); and IMWITOR 742 (caprylic/capric glycerides). (See U.S. Patent No. 6,267,985).

[0054] Still other commercial surfactant compositions having significant triglyceride content are known to those skilled in the art. It should be appreciated

that such compositions, which contain triglycerides as well as surfactants, may be suitable to provide all or part of the lipophilic phase component of the of the present invention, as well as all or part of the surfactants.

[0055] The pharmaceutical compositions of the present invention also comprise an active vitamin D compound. The term "active vitamin D compound," as used herein, is intended to refer to vitamin D which has been hydroxylated in at least the carbon-1 position of the A ring, *e.g.*, 1 $\alpha$ -hydroxyvitamin D<sub>3</sub>. The preferred active vitamin D compound in relation to the composition of the present invention is 1 $\alpha$ ,25-hydroxyvitamin D<sub>3</sub>, also known as calcitriol. A large number of other active vitamin D compounds are known and can be used in the practice of the invention. Examples include 1 $\alpha$ -hydroxy derivatives with a 17 side chain greater in length than the cholesterol or ergosterol side chains (*see* U.S. Patent No. 4,717,721); cyclopentano-vitamin D analogs (*see* U.S. Patent No. 4,851,401); vitamin D<sub>3</sub> analogues with alkynyl, alkenyl, and alkanyl side chains (*see* U.S. Patent Nos. 4,866,048 and 5,145,846); trihydroxycalciferol (*see* U.S. Patent No. 5,120,722); fluoro-cholecalciferol compounds (*see* U.S. Patent No. 5,547,947); methyl substituted vitamin D (*see* U.S. Patent No. 5,446,035); 23-oxa-derivatives (*see* U.S. Patent No. 5,411,949); 19-nor-vitamin D compounds (*see* U.S. Patent No. 5,237,110); and hydroxylated 24-homo-vitamin D derivatives (*see* U.S. Patent No. 4,857,518). Particular examples include ROCALTROL (Roche Laboratories); CALCIJEX injectable calcitriol; investigational drugs from Leo Pharmaceuticals including EB 1089 (24a,26a,27a-trihomo-22,24-diene-1 $\alpha$ ,25-(OH)<sub>2</sub>-D<sub>3</sub>, KH 1060 (20-epi-22-oxa-24a,26a,27a-trihomo-1 $\alpha$ ,25-(OH)<sub>2</sub>-D<sub>3</sub>), Seocalcitol, MC 1288 (1,25-(OH)<sub>2</sub>-20-epi-D<sub>3</sub>) and MC 903 (calcipotriol, 1 $\alpha$ 24s-(OH)<sub>2</sub>-22-ene-26,27-dehydro-D<sub>3</sub>); Roche Pharmaceutical drugs that include 1,25-(OH)<sub>2</sub>-16-ene-D<sub>3</sub>, 1,25-(OH)<sub>2</sub>-16-ene-23-yne-D<sub>3</sub>, and 25-(OH)<sub>2</sub>-16-ene-23-yne-D<sub>3</sub>; Chugai Pharmaceuticals 22-oxacalcitriol (22-oxa-1 $\alpha$ ,25-(OH)<sub>2</sub>-D<sub>3</sub>; 1 $\alpha$ -(OH)-D<sub>5</sub> from the University of Illinois; and drugs from the Institute of Medical Chemistry-Schering AG that include ZK 161422 (20-methyl-1,25-(OH)<sub>2</sub>-D<sub>3</sub>) and ZK 157202 (20-methyl-23-ene-1,25-(OH)<sub>2</sub>-D<sub>3</sub>); 1 $\alpha$ -(OH)-D<sub>2</sub>; 1 $\alpha$ -(OH)-D<sub>3</sub> and 1 $\alpha$ -(OH)-D<sub>4</sub>. Additional examples include 1 $\alpha$ ,25-(OH)<sub>2</sub>-26,27-d<sub>6</sub>-D<sub>3</sub>; 1 $\alpha$ ,25-(OH)<sub>2</sub>-



22-ene-D<sub>3</sub>; 1 $\alpha$ ,25-(OH)<sub>2</sub>-D<sub>3</sub>; 1 $\alpha$ ,25-(OH)<sub>2</sub>-D<sub>2</sub>; 1 $\alpha$ ,25-(OH)<sub>2</sub>-D<sub>4</sub>; 1 $\alpha$ ,24,25-(OH)<sub>3</sub>-D<sub>3</sub>; 1 $\alpha$ ,24,25-(OH)<sub>3</sub>-D<sub>2</sub>; 1 $\alpha$ ,24,25-(OH)<sub>3</sub>-D<sub>4</sub>; 1 $\alpha$ -(OH)-25-FD<sub>3</sub>; 1 $\alpha$ -(OH)-25-FD<sub>4</sub>; 1 $\alpha$ -(OH)-25-FD<sub>2</sub>; 1 $\alpha$ ,24-(OH)<sub>2</sub>-D<sub>4</sub>; 1 $\alpha$ ,24-(OH)<sub>2</sub>-D<sub>3</sub>; 1 $\alpha$ ,24-(OH)<sub>2</sub>-D<sub>2</sub>; 1 $\alpha$ ,24-(OH)<sub>2</sub>-25-FD<sub>4</sub>; 1 $\alpha$ ,24-(OH)<sub>2</sub>-25-FD<sub>3</sub>; 1 $\alpha$ ,24-(OH)<sub>2</sub>-25-FD<sub>2</sub>; 1 $\alpha$ ,25-(OH)<sub>2</sub>-26,27-F<sub>6</sub>-22-ene-D<sub>3</sub>; 1 $\alpha$ ,25-(OH)<sub>2</sub>-26,27-F<sub>6</sub>-D<sub>3</sub>; 1 $\alpha$ ,25S-(OH)<sub>2</sub>-26-F<sub>3</sub>-D<sub>3</sub>; 1 $\alpha$ ,25-(OH)<sub>2</sub>-24-F<sub>1</sub>-D<sub>3</sub>; 1 $\alpha$ ,25S,26-(OH)<sub>2</sub>-22-ene-D<sub>3</sub>; 1 $\alpha$ ,25R,26-(OH)<sub>2</sub>-22-ene-D<sub>3</sub>; 1 $\alpha$ ,25-(OH)<sub>2</sub>-D<sub>2</sub>; 1 $\alpha$ ,25-(OH)<sub>2</sub>-24-epi-D<sub>3</sub>; 1 $\alpha$ ,25-(OH)<sub>2</sub>-23-yne-D<sub>3</sub>; 1 $\alpha$ ,25-(OH)<sub>2</sub>-24R-F-D<sub>3</sub>; 1 $\alpha$ ,25S,26-(OH)<sub>2</sub>-D<sub>3</sub>; 1 $\alpha$ ,24R-(OH)<sub>2</sub>-25F-D<sub>3</sub>; 1 $\alpha$ ,25-(OH)<sub>2</sub>-26,27-F<sub>6</sub>-23-yne-D<sub>3</sub>; 1 $\alpha$ ,25R-(OH)<sub>2</sub>-26-F<sub>3</sub>-D<sub>3</sub>; 1 $\alpha$ ,25,28-(OH)<sub>3</sub>-D<sub>2</sub>; 1 $\alpha$ ,25-(OH)<sub>2</sub>-16-ene-23-yne-D<sub>3</sub>; 1 $\alpha$ ,24R,25-(OH)<sub>3</sub>-D<sub>3</sub>; 1 $\alpha$ ,25-(OH)<sub>2</sub>-26,27-F<sub>6</sub>-23-ene-D<sub>3</sub>; 1 $\alpha$ ,25R-(OH)<sub>2</sub>-22-ene-26-F<sub>3</sub>-D<sub>3</sub>; 1 $\alpha$ ,25S-(OH)<sub>2</sub>-22-ene-26-F<sub>3</sub>-D<sub>3</sub>; 1 $\alpha$ ,25R-(OH)<sub>2</sub>-D<sub>3</sub>-26,26,26-d<sub>3</sub>; 1 $\alpha$ ,25S-(OH)<sub>2</sub>-D<sub>3</sub>-26,26,26-d<sub>3</sub>; and 1 $\alpha$ ,25R-(OH)<sub>2</sub>-22-ene-D<sub>3</sub>-26,26,26-d<sub>3</sub>. Additional examples can be found in WO 99/49870. *See also, e.g.,* U.S. Patent Nos. 5,457,217, 5,447,924, 5,446,034, 5,414,098, 5,403,940, 5,384,313, 5,374,629, 5,373,004, 5,371,249, 5,430,196, 5,260,290, 5,393,749, 5,395,830, 5,250,523, 5,247,104, 5,397,775, 5,194,431, 5,281,731, 5,254,538, 5,232,836, 5,185,150, 5,321,018, 5,086,191, 5,036,061, 5,030,772, 5,246,925, 4,973,584, 5,354,744, 4,927,815, 4,804,502, 4,857,518, 4,851,401, 4,851,400, 4,847,012, 4,755,329, 4,940,700, 4,619,920, 4,594,192, 4,588,716, 4,564,474, 4,552,698, 4,588,528, 4,719,204, 4,719,205, 4,689,180, 4,505,906, 4,769,181, 4,502,991, 4,481,198, 4,448,726, 4,448,721, 4,428,946, 4,411,833, 4,367,177, 4,336,193, 4,360,472, 4,360,471, 4,307,231, 4,307,025, 4,358,406, 4,305,880, 4,279,826, and 4,248,791.

[0056] The pharmaceutical compositions of the present invention may further comprise one or more additives. Additives that are well known in the art include, *e.g.*, detackifiers, anti-foaming agents, buffering agents, antioxidants (*e.g.*, ascorbyl palmitate, butyl hydroxy anisole (BHA), butyl hydroxy toluene (BHT) and tocopherols, *e.g.*,  $\alpha$ -tocopherol (vitamin E)), preservatives, chelating agents, viscomodulators, tonicifiers, flavorants, colorants, odorants, opacifiers, suspending agents, binders, fillers, plasticizers, lubricants, and mixtures thereof.

The amounts of such additives can be readily determined by one skilled in the art, according to the particular properties desired.

[0057] The additive may also comprise a thickening agent. Suitable thickening agents may be of those known and employed in the art, including, *e.g.*, pharmaceutically acceptable polymeric materials and inorganic thickening agents. Exemplary thickening agents for use in the present pharmaceutical compositions include polyacrylate and polyacrylate co-polymer resins, for example poly-acrylic acid and poly-acrylic acid/methacrylic acid resins; celluloses and cellulose derivatives including: alkyl celluloses, *e.g.*, methyl-, ethyl- and propyl-celluloses; hydroxyalkyl-celluloses, *e.g.*, hydroxypropyl-celluloses and hydroxypropylalkyl-celluloses such as hydroxypropyl-methyl-celluloses; acylated celluloses, *e.g.*, cellulose-acetates, cellulose-acetatephthallates, cellulose-acetatesuccinates and hydroxypropylmethyl-cellulose phthallates; and salts thereof such as sodium-carboxymethyl-celluloses; polyvinylpyrrolidones, including for example poly-N-vinylpyrrolidones and vinylpyrrolidone co-polymers such as vinylpyrrolidone-vinylacetate co-polymers; polyvinyl resins, *e.g.*, including polyvinylacetates and alcohols, as well as other polymeric materials including gum traganth, gum arabicum, alginates, *e.g.*, alginic acid, and salts thereof, *e.g.*, sodium alginates; and inorganic thickening agents such as atapulgite, bentonite and silicates including hydrophilic silicon dioxide products, *e.g.*, alkylated (for example methylated) silica gels, in particular colloidal silicon dioxide products.

[0058] Such thickening agents as described above may be included, *e.g.*, to provide a sustained release effect. However, where oral administration is intended, the use of thickening agents as aforesaid will generally not be required and is generally less preferred. Use of thickening agents is, on the other hand, indicated, *e.g.*, where topical application is foreseen.

[0059] Compositions in accordance with the present invention may be employed for administration in any appropriate manner, *e.g.*, orally, *e.g.*, in unit dosage form, for example in a solution, in hard or soft encapsulated form including gelatin encapsulated form, *e.g.*, parenterally or topically, *e.g.*, for application to the skin, for example in the form of a cream, paste, lotion, gel, ointment, poultice,

cataplasm, plaster, dermal patch or the like, or for ophthalmic application, for example in the form of an eye-drop, -lotion or -gel formulation. Readily flowable forms, for example solutions and emulsions, may also be employed *e.g.*, for intralesional injection, or may be administered rectally, *e.g.*, as an enema.

[0060] When the composition of the present invention is formulated in unit dosage form, the active vitamin D compound will preferably be present in an amount of between 10 and 75  $\mu\text{g}$  per unit dose. More preferably, the amount of active vitamin D compound per unit dose will be about 10  $\mu\text{g}$ , 15  $\mu\text{g}$ , 20  $\mu\text{g}$ , 25  $\mu\text{g}$ , 30  $\mu\text{g}$ , 35  $\mu\text{g}$ , 40  $\mu\text{g}$ , 45  $\mu\text{g}$ , 50  $\mu\text{g}$ , 55  $\mu\text{g}$ , 60  $\mu\text{g}$ , 65  $\mu\text{g}$ , 70  $\mu\text{g}$ , or 75  $\mu\text{g}$ .

[0061] When the unit dosage form of the composition is a capsule, the total quantity of ingredients present in the capsule is preferably about 10-1000  $\mu\text{L}$ . More preferably, the total quantity of ingredients present in the capsule is about 100-300  $\mu\text{L}$ .

[0062] The relative proportion of ingredients in the compositions of the invention will, of course, vary considerably depending on the particular type of composition concerned. The relative proportions will also vary depending on the particular function of ingredients in the composition. The relative proportions will also vary depending on the particular ingredients employed and the desired physical characteristics of the product composition, *e.g.*, in the case of a composition for topical use, whether this is to be a free flowing liquid or a paste. Determination of workable proportions in any particular instance will generally be within the capability of a person of ordinary skill in the art. All indicated proportions and relative weight ranges described below are accordingly to be understood as being indicative of preferred or individually inventive teachings only and not as not limiting the invention in its broadest aspect.

[0063] The lipophilic phase component of the invention will suitably be present in an amount of from about 30% to about 90% by weight based upon the total weight of the composition. Preferably, the lipophilic phase component is present in an amount of from about 50% to about 85% by weight based upon the total weight of the composition.

[0064] The surfactant or surfactants of the invention will suitably be present in an amount of from about 1% to 50% by weight based upon the total weight of the composition. Preferably, the surfactant(s) is present in an amount of from about 5% to about 40% by weight based upon the total weight of the composition.

[0065] The amount of active vitamin D compound in compositions of the invention will of course vary, *e.g.*, depending on the intended route of administration and to what extent other components are present. In general, however, the active vitamin D compound of the invention will suitably be present in an amount of from about 0.005% to 20% by weight based upon the total weight of the composition. Preferably, the active vitamin D compound is present in an amount of from about 0.01% to 15% by weight based upon the total weight of the composition.

[0066] The hydrophilic phase component of the invention will suitably be present in an amount of from about 2% to about 20% by weight based upon the total weight of the composition. Preferably, the hydrophilic phase component is present in an amount of from about 5% to 15% by weight based upon the total weight of the composition.

[0067] The pharmaceutical composition of the invention may be in a semisolid formulation. Semisolid formulations within the scope of the invention may comprise, *e.g.*, a lipophilic phase component present in an amount of from about 60% to about 80% by weight based upon the total weight of the composition, a surfactant present in an amount of from about 5% to about 35% by weight based upon the total weight of the composition, and an active vitamin D compound present in an amount of from about 0.01% to about 15% by weight based upon the total weight of the composition.

[0068] The pharmaceutical compositions of the invention may be in a liquid formulation. Liquid formulations within the scope of the invention may comprise, *e.g.*, a lipophilic phase component present in an amount of from about 50% to about 60% by weight based upon the total weight of the composition, a surfactant present in an amount of from about 4% to about 25% by weight based upon the total weight of the composition, an active vitamin D compound present

in an amount of from about 0.01% to about 15% by weight based upon the total weight of the composition, and a hydrophilic phase component present in an amount of from about 5% to about 10% by weight based upon the total weight of the composition.

[0069] In addition to the foregoing the present invention also provides a process for the production of a pharmaceutical composition as hereinbefore defined, which process comprises bringing the individual components thereof into intimate admixture and, when required, compounding the obtained composition in unit dosage form, for example filling said composition into gelatin, *e.g.*, soft or hard gelatin, capsules, or non-gelatin capsules.

[0070] In a more particular embodiment, the invention provides a process for the preparation of a pharmaceutical composition, which process comprises bringing an active vitamin D compound, *e.g.*, calcitriol, into close admixture with a lipophilic phase component and a surfactant as hereinbefore defined, the relative proportion of the lipophilic phase component and the surfactant being selected relative to the quantity of active vitamin D compound employed, such that an emulsion pre-concentrate is obtained.

[0071] The present invention also provides methods for the treatment and prevention of hyperproliferative diseases such as cancer and psoriasis, said methods comprising administering an active vitamin D compound in an emulsion pre-concentrate formulation to a patient in need thereof. Alternatively, the active vitamin D compound can be administered in an emulsion formulation that is made by diluting an emulsion pre-concentrate of the present invention with an appropriate quantity of water.

[0072] Cancers which can be treated with the formulations of the invention include any cancer treatable by an active vitamin D compound. Such cancers include without limitation cancers of the prostate, breast, colon, lung, head and neck, pancreas, endometrium, bladder, cervix, ovaries, squamous cell carcinoma, renal cell carcinoma, myeloid and lymphocytic leukemia, lymphoma, medullary thyroid carcinoma, melanoma, multiple myeloma, retinoblastoma and sarcomas of the soft tissues and bone.

[0073] Preferably, the cancers are treated according to the pulse dose protocols disclosed in WO 99/49870. In this embodiment, the formulations are administered no more than once every three days, more preferably, no more than once a week, more preferably, no more than once every ten days. Preferably, about 5 to about 100  $\mu\text{g}$  of calcitriol, more preferably, about 10 to 60  $\mu\text{g}$ , more preferably, about 40-50  $\mu\text{g}$  of calcitriol, or an equivalent amount of another active vitamin D compound, is administered to an animal in need thereof.

[0074] Animals which may be treated according to the present invention include all animals which may benefit from administration of the formulations of the present invention. Such animals include humans, pets such as dogs and cats, and veterinary animals such as cows, pigs, sheep, goats and the like.

[0075] The following examples are illustrative, but not limiting, of the method and compositions of the present invention. Other suitable modifications and adaptations of the variety of conditions and parameters normally encountered in clinical therapy and which are obvious to those skilled in the art are within the spirit and scope of the invention.

## EXAMPLES

### EXAMPLE 1

#### Relative Chemical Compatibility of Calcitriol With Selected Components

[0076] In this example, the relative chemical compatibility of calcitriol with selected lipophilic, hydrophilic and surfactant components was evaluated by measuring the percent recovery of intact calcitriol after storage at 40°C and 60°C. Calcitriol recovery was determined based on analyses of high-pressure liquid chromatography (HPLC). The results are presented in Table 1.

-22-

TABLE 1: Percent Recovery of Calcitriol Formulated in  
Selected Components

Component	Excipient	Time	% Recovery at 40°C	% Recovery at 60°C
Lipophilic	Corn oil	0	100.00	100.00
		3 days	93.77	104.80
		7 days	90.27	91.50
		14 days	89.89	86.46
	Soybean oil	0	100.00	100.00
		3 days	96.44	94.56
		7 days	98.46	98.57
		14 days	96.66	93.15
	Sunflower oil	0	100.00	100.00
		3 days	99.10	99.33
		7 days	102.77	102.93
		14 days	96.56	88.79
	Vitamin E	0	100.00	100.00
		3 days	128.56	160.79
		7 days	0.00	0.00
		14 days	102.29	65.02
	Miglyol 812	0	100.00	100.00
		3 days	98.23	97.01
		7 days	99.31	96.78
		14 days	99.17	99.48
	Miglyol 812, 0.02% BHA/BHT	0	100.00	100.00
		3 days	98.41	97.83
		7 days	97.43	98.17
		14 days	98.72	102.15
	Captex 200	0	100.00	100.00
		3 days	99.20	97.28
		7 days	100.14	97.68
		14 days	108.83	101.15
	Labrafac CC	0	100.00	100.00
		3 days	98.60	95.84
		7 days	100.05	99.51
		14 days	101.37	100.24

-23-

Component	Excipient	Time	% Recovery at 40°C	% Recovery at 60°C
Hydrophilic	PEG 300	0	100.00	100.00
		3 days	78.22	18.95
		7 days	52.68	4.61
		14 days	10.09	1.84
	Propylene Glycol	0	100.00	100.00
		3 days	97.56	99.71
		7 days	101.73	108.47
		14 days	105.83	138.22
	Cremophor ELP	0	100.00	100.00
		3 days	82.61	66.28
		7 days	62.86	60.90
		14 days	51.90	59.92
Surfactant	Cremophor RH 40 25%	0	100.00	100.00
		3 days	105.30	91.91
		7 days	92.10	78.30
		14 days	96.88	87.95
	in Miglyol 812			
	Polysorbate 80	0	100.00	100.00
		3 days	87.94	67.43
		7 days	87.29	71.71
		14 days	60.52	66.08
	GELUCIRE 44/14 25%	0	100.00	100.00
		3 days	98.70	107.68
		7 days	101.55	83.06
		14 days	100.96	98.11
	in Miglyol 812			
	Vitamin E TPGS 25%	0	100.00	100.00
		3 days	101.15	97.26
		7 days	101.26	98.74
		14 days	103.61	100.15
	in Miglyol 812			
	Labrifil M	0	100.00	100.00
		3 days	98.46	95.19
		7 days	99.45	95.64
		14 days	100.30	78.97
	Poloxamer 188 25%	0	100.00	100.00
		3 days	116.42	76.47
		7 days	126.39	116.67
		14 days	126.79	83.30
	in Miglyol 812			



[0077] The recovery data suggest that the most compatible components are Miglyol 812 (with or without BHT and BHA), Labrafac CC and Captex 200 in the lipophilic component group, propylene glycol in the hydrophilic group, and vitamin E TPGS and GELUCIRE 44/14 in the surfactant group.

## EXAMPLE 2

### Stability of Liquid and Semi-Solid Calcitriol Formulations

#### I. Introduction

[0078] In this Example, the stability of the active vitamin D compound calcitriol was measured in nine different formulations (four liquid formulations and five semisolid formulations).

#### II. Preparation of Calcitriol Formulations

##### A. Liquid Formulations

[0079] Four liquid calcitriol formulations (L1-L4) were prepared containing the ingredients listed in Table 2. The final formulation contains 0.208 mg calcitriol per gram of liquid formulation.

TABLE 2: Composition of Liquid Calcitriol Formulations

Ingredient	L1	L2	L3	L4
Calcitriol	0.0208	0.0208	0.0208	0.0208
Miglyol 812	56.0	62.0	0	0
Captex 200	0	0	55.0	0
Labrafac CC	0	0	0	55.0
Vitamin-E TPGS	15.0	24.0	22.0	20.0
Labrifil M	23.0	4.0	14.0	15.0
1,2-propylene glycol	6.0	10.0	9.0	10.0
BHT	0.05	0.05	0.05	0.05
BHA	0.05	0.05	0.05	0.05

Amounts shown are in grams.

#### B. Semi-Solid Formulations

[0080] Five semi-solid calcitriol formulations (SS1-SS5) were prepared containing the ingredients listed in Table 3. The final formulation contains 0.208 mg calcitriol per gram of semi-solid formulation.

TABLE 3: Composition of Semi-Solid Calcitriol Formulations

Ingredient	SS1	SS2	SS3	SS4	SS5
Calcitriol	0.0208	0.0208	0.0208	0.0208	0.0208
Miglyol 812	80.0	0	65.0	0	79.0
Captex 200	0	82.0	0	60.0	0
Labrafac CC	0	0	0	0	12.0
Vitamin-E TPGS	20.0	18.0	5.0	5.0	9.0
Labrifil M	0	0	0	0	0
Gelucire 44/14	0	0	30.0	35.0	0
BHT	0.05	0.05	0.05	0.05	0.05
BHA	0.05	0.05	0.05	0.05	0.05

Amounts shown are in grams.

### C. Method of Making the Liquid and Semi-Solid Calcitriol Formulations

#### 1. Preparation of Vehicles

[0081] One hundred gram quantities of the four liquid calcitriol formulations (L1-L4) and the five semi-solid calcitriol formulations (SS1-SS5) listed in Tables 2 and 3, respectively, were prepared as follows.

[0082] The listed ingredients, except for calcitriol, were combined in a suitable glass container and mixed until homogeneous. Vitamin E TPGS and GELUCIRE 44/14 were heated and homogenized at 60°C prior to weighing and adding into the formulation.

#### 2. Preparation of Active Formulations

[0083] The semi-solid vehicles were heated and homogenized at  $\leq 60^{\circ}\text{C}$ . Under subdued light,  $12 \pm 1$  mg of calcitriol was weighed out into separate glass bottles with screw caps, one bottle for each formulation. (Calcitriol is light-sensitive;

-27-

subdued light/red light should be used when working with calcitriol/calcitriol formulations.) The exact weight was recorded to 0.1 mg. The caps were then placed on the bottles as soon as the calcitriol had been placed into the bottles. Next, the amount of each vehicle required to bring the concentration to 0.208 mg/g was calculated using the following formula:

$$C_w/0.208 = \text{required weight of vehicle}$$

Where  $C_w$  = weight of calcitriol, in mg, and

0.208 = final concentration of calcitriol (mg/g).

[0084] Finally, the appropriate amount of each vehicle was added to the respective bottle containing the calcitriol. The formulations were heated ( $\leq 60^\circ\text{C}$ ) while being mixed to dissolve the calcitriol.

### III. Stability of Calcitriol Formulations

[0085] The nine calcitriol formulations (L1-L4 and SS1-SS5) were analyzed for stability of the calcitriol component at three different temperatures. Sample of the nine formulations were each placed at  $25^\circ\text{C}$ ,  $40^\circ\text{C}$ , and  $60^\circ\text{C}$ . Samples from all three temperatures for all nine formulations were analyzed by HPLC after 1, 2 and 3 weeks. In addition, samples from the  $60^\circ\text{C}$  experiment were analyzed by HPLC after 9 weeks. The percent of the initial calcitriol concentration remaining at each time point was determined for each sample and is reported in Table 4 (liquid formulations) and Table 5 (semi-solid formulations).

TABLE 4: Stability of Liquid Formulations

Formulation	Temp.	Recovery* of Calcitriol (%)			
		Week 1	Week 2	Week 3	Week 9
Liquid #1	25°C	99.3	98.6	99.7	ND
	40°C	103.2	100.4	100.2	ND
	60°C	99.4	98.4	98.4	91.7
Liquid #2	25°C	98.1	95.2	97.7	ND
	40°C	98.0	97.1	99.2	ND
	60°C	97.1	95.6	96.7	93.1
Liquid #3	25°C	99.7	99.2	102.3	ND
	40°C	100.1	99.9	100.7	ND
	60°C	98.3	98.7	98.4	90.5
Liquid #4	25°C	98.4	97.7	98.0	ND
	40°C	100.0	101.0	100.8	ND
	60°C	98.5	97.5	99.0	86.1

\* Percent of time zero concentration.

TABLE 5: Stability of Semi-Solid Formulations

Formulation	Temp.	Recovery* of Calcitriol (%)			
		Week 1	Week 2	Week 3	Week 9
Semi-Solid #1	25°C	98.5	98.9	99.8	ND
	40°C	99.6	99.0	98.2	ND
	60°C	97.9	97.2	96.3	104.6
Semi-Solid #2	25°C	100.0	99.6	100.4	ND
	40°C	98.7	99.6	98.7	ND
	60°C	97.2	98.0	98.6	100.0
Semi-Solid #3	25°C	101.2	98.9	100.4	ND
	40°C	100.0	98.7	98.8	ND
	60°C	98.3	97.6	98.4	97.1
Semi-Solid #4	25°C	100.2	99.0	99.6	ND
	40°C	98.4	99.2	98.5	ND
	60°C	96.8	97.7	97.7	103.4
Semi-Solid #5	25°C	98.8	99.2	98.9	ND
	40°C	99.0	97.1	96.8	ND
	60°C	96.8	96.7	96.0	97.7

\* Percent of time zero concentration.

[0086] As illustrated by Tables 4 and 5, calcitriol remained relatively stable with very little degradation in all of the formulations (liquid and semi-solid) analyzed.

### EXAMPLE 3

#### Appearance and UV/Visible Absorption Study of Calcitriol Formulations

[0087] Calcitriol formulations L1 and SS3 were prepared prior to this study and stored at room temperature protected from light. Table 6 below shows the quantities of ingredients used to prepare the formulations.

TABLE 6: Composition of Calcitriol Formulations Used  
for Absorption Analysis

Ingredient	Liquid #1	Semi-Solid #3
Calcitriol	0.0131	0.0136
Vitamin-E TPGS	9.45	3.27
Miglyol 812	35.28	42.51
Labrifil M	14.49	0
Gelucire 44/14	0	19.62
1,2-propylene glycol	3.78	0
BHA	0.03	0.03
BHT	0.03	0.03

Amounts shown are in grams.

[0088] The formulations were warmed to 55°C prior to use. Both formulations (liquid #1 and semi-solid #3) were mixed well with a vortex mixer and appeared as clear liquids. Each calcitriol formulation ( $\approx 250 \mu\text{L}$ ) was added to a 25 mL volumetric flask. The exact weights added were 249.8 mg for Liquid-1 and 252.6 mg for semi-solid #3. Upon contact with the glass, the semi-solid-3 formulation became solidified. Deionized water was then added to the 25 mL mark and the solutions were mixed with a vortex mixer until uniform. The appearance was observed at this point and the absorbance of the resulting mixtures at 400 nm was determined by UV/visible spectrophotometry. Deionized water was used as a blank and the measurements were taken at 400 nm. Each sample was measured 10 times over a period of 10 minutes. The results are summarized in Table 7. Both formulations formed were white and opaque.

TABLE 7: Absorption Readings of the Formulations at 400 nm

Measurement	Liquid #1	Semi-Solid #3
1	2.4831	1.6253
2	2.5258	1.6290
3	2.5411	1.6309
4	2.5569	1.6328
5	2.5411	1.6328
6	2.5258	1.6347
7	2.5569	1.6328
8	2.5111	1.6366
9	2.5111	1.6366
10	2.5411	1.6328
Average	2.5294	1.6324
RSD%	0.91	0.21

## EXAMPLE 4

## Diameter of Emulsion Droplets Formed From the Liquid and Semi-Solid Formulation Vehicles (without calcitriol)

[0089] In this example, the average diameter of emulsion droplets was measured after dilution of the liquid (L1-L4) and semi-solid (SS1-SS5) emulsion pre-concentrate vehicles (not containing calcitriol) with simulated gastric fluid (SGF) lacking enzyme. The average diameter of the droplets was determined based on light scattering measurements. The appearance of the pre-concentrates and the resulting emulsions, determined by visual inspection, was also noted. The results are summarized in Table 8.



**TABLE 8: Diameter of Emulsion Droplets Formed From Emulsion Pre-Concentrate Vehicles (without calcitriol)**

Formulation	Appearance of emulsion pre-concentrate	pre-concentrate: SGF ratio	Ave. hydro-dynamic diameter*	Appearance of emulsion
L1	Clear liquid	1:1600	237	opaque
L2	Clear liquid	1:1600	281	opaque
L3	Clear liquid	1:1600	175	opaque
L4	Clear liquid	1:1600	273	opaque
SS1	Semi-solid	1:2000	305	opaque
SS2	Semi-solid	1:2000	259	opaque
SS3	Semi-solid	1:2000	243	opaque
SS4	Semi-solid	1:2000	253	opaque
SS5	Semi-solid	1:2000	267	opaque

\*(Zaverage in nanometer)

[0090] From the results presented above, it is concluded that the droplets (particles) formed from the emulsion preconcentrate formulations were of sub-micron droplet size despite having an opaque appearance.

#### EXAMPLE 5

##### Diameter of Emulsion Droplets Formed From Liquid and Semi-Solid Calcitriol Formulation

[0091] In this example, the average diameter of emulsion droplets was measured after dilution of the liquid #1 (L1) and semi-solid #3 (SS3) emulsion pre-concentrates in simulated gastric fluid (SGF) without enzyme. The formulations used in this example contained calcitriol at a concentration of 0.2 mg calcitriol/g of formulation. The diameter of the droplets was determined based on light scattering measurements. The appearance of the resulting emulsions, determined by visual inspection, was also noted. The results are summarized in Table 9.

TABLE 9: Diameter of Emulsion Droplets Formed From Emulsion Pre-Concentrate Formulations Containing Calcitriol

Formulation	pre-concentrate: SGF ratio	Ave. hydro- dynamic diameter*	Appearance of emulsion
L1	1:1600	257	opaque
SS3	1:2000	263	opaque

\*(Zaverage in nanometer)

#### EXAMPLE 6

##### *In Vitro* Dispersion of Calcitriol From Emulsion Pre-Concentrates

[0092] In this Example, the extent of calcitriol dispersion in various formulations in gelatin capsules was determined. A single capsule containing 250 mg of a calcitriol formulation in a size-2 gelatin capsule (each capsule containing 0.2mg calcitriol/g formulation) was added to 200 mL of simulated gastric fluid (SGF) without enzyme at 37°C and was mixed by a paddle at 200 RPM. Samples were then filtered through a 5 µm filter and analyzed for calcitriol concentration at 30, 60, 90, and 120 minutes by HPLC. The results are shown in Table 10.

TABLE 10: Percent Calcitriol Obtained in Filtrate After Dispersion in SGF and Filtration Through a 5 µm Filter

Formulation	30 min.	60 min.	90 min.	120 min.
Liquid #1	106	103	86	68
Semi-Solid #3	109	99	73	53
Comparison Formulation <sup>#</sup>	0	0	0	0

<sup>#</sup>The Comparison Formulation contained calcitriol at 0.2 mg/g dissolved in Miglyol 812 with 0.05% BHA and 0.05% BHT. This formulation is similar to the ROCALTROL formulation available from Roche Laboratories.

[0093] As this Example illustrates, the dispersion of calcitriol in simulated gastric fluid from capsules containing either the L1 or the SS3 formulations was much more extensive than that which was observed with capsules containing the

Comparison Formulation (which is similar to the ROCALTROL formulation available from Roche Laboratories).

#### EXAMPLE 7

##### Plasma Concentrations and Pharmacokinetics of Calcitriol in Dogs

[0094] A pharmacokinetics study in dogs compared the plasma levels of calcitriol after administration of 1.0 µg/kg using 3 different formulations: ROCALTROL, a liquid formulation (liquid #1, and a semi-solid formulation (semi-solid #3). Four dogs received 1.0 µg/kg orally of ROCALTROL, the semi-solid formulation, or the liquid formulation. When dogs were used for more than one formulation a minimum 7-day washout period separated dosing with each formulation.

[0095] Blood samples were obtained pre-dose, and 0.5, 1, 2, 4, 6, 8, 10, 12, 24, 36, and 48 hours post-dose for analysis of calcitriol levels. Blood samples for clinical chemistry were obtained pre-dose, and at 24 and 48 hours post-dose for the ROCALTROL group; samples were obtained pre-dose, and at 4, 24, 48, 72, 96, and 120 hours for the semi-solid and liquid formulations. Samples were analyzed for calcitriol by radioimmunoassay and subjected to pharmacokinetics analyses.

[0096] Plasma concentrations of calcitriol over time for the three formulations are shown graphically in Figure 1.

[0097] A summary of the pharmacokinetics of calcitriol as one of three different formulations at a common dose of 1.0 µg/kg is presented in Tables 11-14.

TABLE 11: Summary of Calcitriol Parameters in Dogs

Parameter	ROCALTROL		Semi-Solid #3		Liquid #1	
	Mean	SD	Mean	SD	Mean	SD
$C_{max}$ , pg/mL	717.4	51.5	2066.6	552.5	2164.4	253.9
$T_{max}$ <sup>a</sup> , h	3.0	(2-6)	2.0	(1-2)	1.5	(1-2)
$AUC_{(0-\infty)}$ , pg·h/mL	11988.0	3804.7	12351.7	1624.9	14997.4	3531.7
$T_{1/2}$ <sup>b</sup> , h	25.1	11.1	4.8	1.2	7.8	3.5

<sup>a</sup> Expressed as median and range<sup>b</sup> Expressed as harmonic mean and pseudo SD based on jackknife variance

TABLE 12: Plasma Concentration (pg/mL) and Pharmacokinetic Parameters of Calcitriol in Dog Following a Single 1 µg/kg Administration of ROCALTROL

Parameter	Time, h	Dog 101	Dog 102	Dog 103	Dog 104	Mean	SD
	0.0	BQL	BQL	BQL	BQL	0	0
	0.5	488.2	304.8	182.7	BQL	243.9	205.4
	1.0	478.2	634.8	500.7	555.7	542.4	69.7
	2.0	518.2	700.8	749.7	765.7	683.6	113.7
	4.0	494.2	658.8	750.7	745.7	662.4	119.8
	6.0	652.2	566.8	496.7	523.7	559.9	68.0
	8.0	381.2	366.8	418.7	381.7	387.1	22.2
	10.0	313.2	212.8	165.7	158.7	212.6	71.2
	12.0	190.2	186.8	189.7	171.7	184.6	8.7
	24.0	78.2	78.8	69.7	97.7	81.1	11.8
	36.0	63.2	83.8	80.7	67.7	73.9	10.0
	48.0	66.2	47.8	45.7	52.7	53.1	9.2
$C_{max}$ , pg/mL		652.2	700.8	750.7	765.7	717.4	51.5
$T_{max}$ <sup>a</sup> , h		6.0	2.0	4.0	2.0	3.0	(2-6)
$AUC_{(0-\infty)}$ , pg·h/mL		17693.6	10094.5	9976.2	10187.5	11988.0	3804.7
$T_{1/2}$ <sup>b</sup> , h		100.4	18.8	20.2	21.3	25.1	11.1

<sup>a</sup> Expressed as median and range<sup>b</sup> Expressed as harmonic mean and pseudo SD based on jackknife varianceBold type - used to calculate  $\lambda$

TABLE 13: Plasma Concentration (pg/mL) and Pharmacokinetic Parameters of Calcitriol in Dog Following a Single 1 µg/kg Administration of Semi-solid #3 Formulation

Parameter	Time, h	Dog 101	Dog 102	Dog 103	Dog 104	Mean	SD
	0.0	BQL	BQL	BQL	BQL	0	0
	0.5	198.1	11.0	BQL	BQL	52.3	97.4
	1.0	1208.1	2246.0	1128.7	503.4	1271.6	722.0
	2.0	1255.1	2110.0	2269.7	2495.4	2032.6	541.9
	4.0	902.1	1371.0	1095.7	1437.4	1201.6	248.5
	6.0	603.1	1039.0	932.7	1112.4	921.8	224.9
	8.0	815.1	441.0	593.7	848.4	674.6	192.4
	10.0	253.1	489.0	285.7	305.4	333.3	106.0
	12.0	213.1	295.0	184.7	170.4	215.8	55.7
	24.0	50.1	37.0	40.7	29.4	39.3	8.6
	36.0	14.1	BQL	BQL	13.6	6.9	8.0
	48.0	BQL	BQL	BQL	BQL	0.0	0.0
$C_{max}$ , pg/mL		1255.1	2246.0	2269.7	2495.4	2066.6	552.5
$T_{max}$ <sup>a</sup> , h		2.0	1.0	2.0	2.0	2.0	(1-2)
$AUC_{(0-\infty)}$ , pg·h/mL		10333.8	14012.9	11813.8	13246.4	12351.7	1624.9
$T_{1/2}$ <sup>b</sup> , h		6.2	3.8	4.1	5.9	4.8	1.2

<sup>a</sup> Expressed as median and range

<sup>b</sup> Expressed as harmonic mean and pseudo SD based on jackknife variance

Bold type - used to calculate  $\lambda$

TABLE 14: Plasma Concentration (pg/mL) and Pharmacokinetic Parameters of Calcitriol in Dogs Following a Single 1 µg/kg Liquid #1 Formulation

Parameter	Time, h	Dog 105	Dog 106	Dog 107	Dog 108	Mean	SD
	0.0	BQL	BQL	BQL	BQL	0	0
	0.5	BQL	57.6	523.0	350.0	232.7	246.9
	1.0	1283.0	238.6	2266.0	2468.0	1563.9	1024.0
	2.0	2028.0	1895.6	2026.0	2373.0	2080.7	204.5
	4.0	1090.0	892.6	1009.0	1771.0	1190.7	395.3
	6.0	871.0	763.6	730.0	1063.0	856.9	150.0
	8.0	301.0	579.6	374.0	562.0	454.2	138.1
	10.0	421.0	520.6	464.0	517.0	480.7	47.4
	12.0	348.0	290.6	170.0	373.0	295.4	90.4
	24.0	42.0	165.6	62.0	202.0	117.9	78.0
	36.0	49.0	111.6	BQL	79.0	59.9	47.4
	48.0	35.0	15.5	BQL	BQL	12.6	16.6
$C_{max}$ , pg/mL		2028.0	1895.6	2266.0	2468.0	2164.4	253.9
$T_{max}$ <sup>a</sup> , h		2.0	2.0	1.0	1.0	1.5	(1-2)
$AUC_{(0-\infty)}$ , pg·h/mL		13474.4	14296.3	12101.0	20117.7	14997.4	3531.7
$T_{1/2}$ <sup>b</sup> , h		10.6	8.5	5.0	10.1	7.8	3.5

<sup>a</sup> Expressed as median and range

<sup>b</sup> Expressed as harmonic mean and pseudo SD based on jackknife variance

Bold type - used to calculate I

[0098] The results of this study show that there were some differences and similarities in the pharmacokinetics between these particular inventive formulations and ROCALTROL as follows:

- $C_{max}$  was approximately three times higher with the liquid and semi-solid formulations than with the ROCALTROL formulation.
- $C_{max}$  was achieved sooner (1 to 2 hours) with the liquid and semi-solid formulations than with the ROCALTROL formulation (2 to 4 hours).
- The overall systemic exposure ( $AUC_{(0-\infty)}$ ) was comparable with the three formulations, although systemic exposure in the first 24-48 hours was greater with the liquid and semi-solid formulations than with ROCALTROL.

[0099] The foregoing results show that the liquid #1 formulation produces the highest  $C_{max}$  and the largest AUC calcitriol values, followed closely by the semi-solid #3 formulation. The ROCALTROL formulation has the lowest  $C_{max}$  and AUC values. It appears that the liquid #1 and semi-solid #3 formulations were

absorbed much faster and produced higher plasma concentration during the first twelve hours and a faster rate of elimination.

#### EXAMPLE 8

##### Pharmacokinetics of the Semi-Solid #3 Formulation After Escalating Doses

[0100] In this study the pharmacokinetics of the semi-solid formulation after escalating oral doses was studied in dogs. Three male and three female Beagle dogs were dosed orally with single doses of 0.5  $\mu\text{g/kg}$  (all six dogs), 0.1  $\mu\text{g/kg}$  (1 male and 1 female), 5.0  $\mu\text{g/kg}$  (2 males and 2 females), and 10.0  $\mu\text{g/kg}$  (all dogs). After the 10.0  $\mu\text{g/kg}$  dose, 2 dogs per sex were euthanized. The remaining male and female dogs continued on study and received doses of 30.0  $\mu\text{g/kg}$  and 100.0  $\mu\text{g/kg}$ . After each dose the animals were held for a 6-day recovery period.

[0101] Blood samples (approximately 1 mL) were collected from each dog pre-dose and at 0, 2 (in all but the 0.5  $\mu\text{g/kg}$  dose), 4, 8, 24, 48, and 96 hours following dose administration. Samples were analyzed for calcitriol by radioimmunoassay and subjected to pharmacokinetic analyses. Plasma concentrations of calcitriol are shown graphically for males and females in Figs. 2A and 2B.

[0102] After dosing with semi-solid #3, maximum plasma concentrations usually occurred at the two hour sampling timepoint. At doses above 0.1  $\mu\text{g/kg}$ , plasma concentrations appeared to decline at a more rapid rate during the first 8 hours than during the 24 to 96 hour time period.

[0103] At the lowest dose of 0.1  $\mu\text{g/kg}$ , plasma concentrations of calcitriol fell below the limit of quantitation after 24 hours. At 0.5  $\mu\text{g/kg}$  and above, measurable concentrations of calcitriol persisted at the 96 hour sampling timepoint. There did not appear to be any remarkable differences between the male and the female dogs.

[0104] Pharmacokinetic parameters for semi-solid #3 at doses ranging from 0.1 to 100.0  $\mu\text{g/kg}$  are summarized in Table 15.

Table 15: Pharmacokinetics of Calcitriol After Escalating Doses of Calcitriol (Semi-solid #3)

Dose ( $\mu\text{g/kg}$ )	0.1		0.5		5.0	
Gender	Male	Female	Male	Female	Male	Female
N	1	1	3	3	2	2
$C_{\max}$ (pg/mL)	566	473	1257	1431	17753	18346
Tmax (hr)	2.0	2.0	4.0	4.0	2.0	2.0
AUC <sub>0-24</sub> (pg·hr/mL)	4311	2654	11431	15598	104,027	107,452
AUC <sub>0-48</sub> (pg·hr/mL)	4311	2654	13584	19330	125,408	126,746
AUC <sub>0-∞</sub> (pg·hr/mL)	4916	2718	15062	21644	200,283	160,681
T <sub>1/2</sub> (hr)	4.2	2.7	17.1	14.2	67.6	36.8

Dose ( $\mu\text{g/kg}$ )	10.0		30.0		100.0	
Gender	Male	Female	Male	Female	Male	Female
N	3	3	1	1	1	1
$C_{\max}$ (pg/mL)	23858	32336	53005	115,896	238,619	211,631
Tmax (hr)	2.7	2.0	2.0	2.0	2.0	2.0
AUC <sub>0-24</sub> (pg·hr/mL)	183,981	203,857	311,841	567,717	1,165,988	1,089,831
AUC <sub>0-48</sub> (pg·hr/mL)	223,977	240,483	370,713	641,469	1,381,424	1,256,007
AUC <sub>0-∞</sub> (pg·hr/mL)	388,600	345,936	531,303	854,841	1,874,997	1,731,873
T <sub>1/2</sub> (hr)	77.7	56.0	56.3	58.2	45.3	53.7

[0105] These pharmacokinetic results indicate the following:

- The systemic exposure of calcitriol appeared to be fairly linear throughout the tested dose range of 0.1 to 100.0  $\mu\text{g/kg}$ . No saturation of absorption was observed.
- The half-life of calcitriol appeared to be dose-dependent. Formulations having a half life of greater than 24 hours are less suitable for high dose pulse administration.



- Weekly dosing with semi-solid #3 at 5.0 µg/kg and above resulted in some accumulation in the plasma. Accumulation was not consistently observed at the lower doses of 0.1 and 0.5 µg/kg.

### EXAMPLE 9

#### A 28 Day Oral Toxicity Study in Dogs with Semi-Solid #3

[0106] In this study a 28-day repeated dose toxicology study of semi-solid #3 was conducted in dogs to assess the pharmacokinetics of calcitriol after weekly oral capsule dosing. Semi-solid #3 or control article capsules were administered on study days 0, 7, 14, 21, and 28. Twelve dogs (6 male, 6 female) received vehicle control (group 1), eight dogs (4 male, 4 female) received 0.1 µg/kg semi-solid #3 (group 2), and eight dogs (4 male, 4 female) received 1.0 µg/kg semi-solid #3 (group 3). Twelve dogs (6 male, 6 female) received 30.0 µg/kg semi-solid #3 on day 0 (group 4). Due to the severity of the clinical response observed after the first 30 µg/kg dose on day 0, dose levels were reduced in this group to 10 µg/kg (males on days 7, 14, 21, and 28) or 5 µg/kg (females on days 7, 14, 21, and 28). Blood samples were collected on each dog pre-dose and at 1, 2, 4, 6, 8, 24, and 48 hours following dosing on study days 0 (first dose) and 21 (fourth weekly dose). All animals were sacrificed on study day 29.

[0107] The pharmacokinetic results for plasma calcitriol for groups 2-4 are summarized in Table 16.

Table 16: Mean Toxicokinetic Parameters of Calcitriol After Weekly Dosing with Semi-Solid #3 in Dogs

DAY 0						
Dose	0.1 µg/kg (Group 2)		1.0 µg/kg (Group 3)		30.0 µg/kg (Group 4)	
Sex (No. of Dogs)	Male (4)	Female (4)	Male (4)	Female (4)	Male (6)	Female (6)
$C_{max}$ , pg/mL	198.7	430.8	2385.0	3419.1	84909.1	57133.3
$T_{max}^a$ , h	1.0	2.0	1.0	1.5	2.0	2.0
$AUC_{0-24}$ , pg·hr/mL	1840.6	3093.4	17144.2	23259.7	496044.6	323573.1
$AUC_{0-48}$ , pg·hr/mL	2130.8	3093.4	19141.6	25794.5	644064.2	365340.7

DAY 24 (Fourth Weekly Dose)						
Dose	0.1 µg/kg (Group 2)		1.0 µg/kg (Group 3)		10.0 µg/kg (Group 4)	5.0 µg/kg (Group 4)
Sex (No. of Dogs)	Male (4)	Female (4)	Male (4)	Female (4)	Male (6)	Female (6)
Dose	0.1	0.1	1.0	1.0	10.0 <sup>b</sup>	5.0 <sup>b</sup>
$C_{max}$ , pg/mL	217.6	398.3	2272.1	2188.6	29061.8	8670.7
$T_{max}^a$ , h	1.0	2.0	1.5	2.0	1.0	2.0
$AUC_{0-24}$ , pg·hr/mL	1956.2	3283.0	19765.4	12947.3	173597.2	46878.1
$AUC_{0-48}$ , pg·hr/mL	2225.9	3640.7	24606.9	15380.0	209732.1	54976.1

<sup>a</sup>The values for  $T_{max}$  are the median values for this parameter. All other parameters shown are mean values.

<sup>b</sup>Doses of semi-solid #3 were lowered beginning on Study Day 7.

Data from the vehicle control dogs (Group 1) were not subjected to pharmacokinetic analysis.

[0108] Figs. 3A and 3B show the adjusted plasma concentration-time curve for calcitriol after oral capsule dosing with semi-solid #3 on study days 0 and 21 in male (Fig. 3A) and female (Fig. 3B) Beagle dogs. Calcitriol values at time 0 on day 0 were subtracted from all subsequent timepoints to adjust for endogenous (baseline) plasma calcitriol

[0109] The results of the study indicate that following:

-42-

- After oral capsule dosing with semi-solid #3, plasma concentrations of calcitriol rose fairly rapidly, reaching peak plasma concentrations within two hours.
- Plasma concentrations of calcitriol decreased at a more rapid rate during the first 8 hours post-dosing than during the later timepoints (24-48 hours), possibly indicating redistribution of calcitriol to extravascular spaces, with subsequent slow release of calcitriol back into the vascular spaces. This observation was more apparent at the higher dose levels than at the lower dose levels.
- At 24 hours post-dosing, plasma concentration of calcitriol had declined to near-baseline values at the low dose of 0.1 µg/kg. However, at the higher doses of calcitriol, dose-related residual concentrations of calcitriol were still evident at the last sampling timepoint (48 hours), although all values returned to pre-dose (baseline) values by one week post-dosing.
- Values for  $C_{max}$  and AUC were fairly proportional to dose throughout the dose range tested (0.1-30.0 µg/kg).
- Values for  $AUC_{0-24}$  at the low dose, which was the no observable adverse effect level (0.1 µg/kg) ranged from 1840.6 - 3283.0 pg·hr/mL.
- Values for  $AUC_{0-24}$  at the mid dose, which was the maximum tolerated dose (1.0 µg/kg) ranged from 12,947.3 – 23,259.7 pg·hr/mL.
- Values for  $AUC_{0-24}$  at doses associated with weight loss and moderate signs of toxicity, ranged from 46,878.1 pg·hr/mL (5.0 µg/kg; females) to 173,597.2 pg·hr/mL (10.0 µg/kg; males).
- Values for  $AUC_{0-24}$  at a dose associated with mortality (30.0 µg/kg) ranged from 323,573.1 – 496,044.6 pg·hr/mL.
- There were no consistent sex differences in any pharmacokinetic parameter.

[0110] Overall, the animals appeared to handle calcitriol similarly after the first dose and after repeated once-weekly dosing, with a few exceptions such as higher values for  $C_{max}$  and AUC on Day 0 compared to Day 21 in the 1.0 µg/kg females (not evident in the males).

## EXAMPLE 10

## Acute Toxicity Study of Three Different Formulations

[0111] In the study described in Example 7, several in-life parameters, including clinical chemistry parameters, were monitored to assess the toxicity of the calcitriol formulations. Blood samples were analyzed for calcium, phosphorus, blood urea nitrogen (BUN), glucose, albumin, bilirubin (total), aspartate aminotransferase (AST), alanine aminotransferase (ALT), alkaline phosphatase (AP), and creatinine.

[0112] No clinical toxicity was seen in any dog with any of the three formulations.

[0113] Hypercalcemia was seen after dosing with 1.0 µg/kg with all three formulations. The group mean and the individual range of serum calcium levels of each of the three different formulations are presented in Table 17.

Table 17: Group Mean Serum Calcium Levels (mg/dL)

Historical Control		ROCALTROL, 1.0 µg/kg						
		0 hr	4 hr	24 hr	48 hr	72 hr	96 hr	120 hr
9.25-11.3 <sup>a</sup> (10.44) <sup>b</sup>	Mean	11.1	NA	13.8*	12.9*	NA	NA	NA
	SD	0.31	NA	0.83	0.26	NA	NA	NA
	Range	10.8-11.5	NA	13.2-15.0	12.6-13.1	NA	NA	NA
Calcitriol, liquid, 1.0 µg/kg								
9.25-11.3 (10.44)	Mean	10.4	10.5	16.1*	14.3*	12.7*	12.5*	12.0*
	SD	0.17	0.37	1.47	1.34	0.53	0.78	0.80
	Range	10.2-10.5	10.1-10.9	13.9-17.0	12.9-15.7	12.0-13.3	11.5-13.4	11.2-13.1
Calcitriol, semi-solid, 1.0 µg/kg								
9.25-11.3 (10.44)	Mean	10.1	10.6	14.3*	14.2*	12.3*	12.6*	12.7*
	SD	0.33	0.29	1.72	1.52	1.35	0.76	0.47
	Range	9.7-10.5	10.7-10.8	12.2-16.4	12.1-15.5	10.8-13.6	11.5-13.1	12.0-13.0

<sup>a</sup> Historical range

<sup>b</sup> Historical mean

\* Mean outside historical range

NA = not available (serum sample not taken)

[0114] In addition to elevations of calcium, elevations of ALT, AST, BUN, and creatinine were observed in all groups.

[0115] In summary, the results of this study indicated that:

- No treatment-related clinical signs were evident in any dog after dosing with any of the formulations (ROCALtrol, liquid, or semi-solid).
- Hypercalcemia at 1.0 µg/kg PO was seen in dogs with all three formulations.
- Time course of the hypercalcemia was comparable among all three formulations up to 48 hours; sampling for the ROCALtrol group did not extend beyond 48 hours.
- Severity of the hypercalcemia was comparable among the three formulations; the highest serum calcium (17.0 mg/dL) occurred at 24 hours in dogs receiving the liquid formulation.
- Mean values for ALT, AST, BUN, and creatinine were observed to be outside the historical range in all treatment groups at one or more timepoints.
- Elevations for BUN and creatinine were greater in the liquid or semi-solid groups; in the absence of a concurrent control group, the significance of this observation is unclear.

#### EXAMPLE 11

##### Acute Maximum Tolerated Dose Study

[0116] In the study described above in Example 8, the acute toxicity and hypercalcemia effects of semi-solid #3 were also assessed to estimate the maximum tolerated dose and to provide data for dose selection of future studies.

[0117] Calcium levels were increased in a dose-related manner at all dose levels in males (Fig. 4A) and females (Fig. 4B). Serum calcium data for the 0.001 and 1.0 µg/kg dose was obtained in male dogs in the study describe in Example 10, and is included here for completeness.

[0118] In summary, this study of semi-solid #3 administered orally via a capsule to male and female Beagle dogs at 0.1, 0.5, 5.0, 10.0, 30.0, and 100.0 µg/kg showed:

- Dose dependent hypercalcemia was the most common laboratory abnormality.
- Elevations of creatinine, urea nitrogen, cholesterol, erythrocytes, hemoglobin, hematocrit, and neutrophils, and a decrease in lymphocytes were seen at doses of 5.0 µg/kg or higher.
- Body weights and food consumption decreased markedly after receiving the 30.0 and 100.0 µg/kg doses; after 100.0 µg/kg, dogs had a noticeable thin appearance and obvious decreased activity.

[0119] Based on these results, the maximum tolerated dose of semi-solid #3 in dogs appeared to be 5.0 µg/kg.

#### EXAMPLE 12

##### A 28 Day Repeated Dose Toxicity Study

[0120] In the study described above in Example 9, the dogs were also assessed for potential toxicity of the semi-solid #3 formulation when administered to dogs by the oral (capsule) route once every seven days for 28 days. The study included assessments of clinical signs, body weights, food consumption, toxicokinetics, clinical pathology including biochemistry, hematology, coagulation, and urinalysis, ophthalmology, cardiology, gross necropsy, organ weight, and full histopathology on all animals. The study design is summarized in Table 18.

Table 18: Study Design for 28-Day Repeated Dose Study in Dogs

Group	No. of Main (Recovery) Animals		Dose Materials	Bulk Dose Level (mg/kg/dose)*	Calcitriol Dose Level (µg/kg/dose)
	Males	Females			
1	4 (2)	4 (2)	Control Article	300**	0
2	4	4	Test Article*	1	0.1
3	4	4	Test Article*	10	1
4	4 (2)	4 (2)	Test Article*	300/100 (males)** 300/50 (females)**	30/10 (Males)** 30/5 (Females)**

\* The test article (calcitriol semi-solid #3) is a formulation containing 0.1 mg of calcitriol per gram.

\*\* Dose reduced to 10 µg/kg in males and 5 µg/kg in females at Week 2; all surviving animals were sacrificed on Day 29.

[0121] Four of the group 4 animals (1 male and 3 females) died or were euthanized moribund during the first three days of the study. No deaths occurred following reduction of the dose level on day 7; there were no deaths in groups 1, 2 or 3.

[0122] In the group 4 animals that died, the most notable clinical abnormalities preceding death primarily included red vomitus, few/no feces, soft stools containing red material, red nasal discharge, shallow/rapid breathing, decreased activity and lateral recumbency.

[0123] Dose-related body weight loss, decreased weight gain, and decreased food consumption were observed in group 3 and 4 animals; group 3 animals were ~11-12% below controls; group 4 animals were 17-24% below controls. No effects on weight gain or food consumption were apparent in group 2 animals.

[0124] There was a trend towards an increase in several RBC and WBC parameters in the group 4 animals at day 29; no toxicologically significant hematological abnormalities were apparent in the group 2 and 3 animals.

[0125] Dose related hypercalcemia was noted in group 3 and 4 animals. Calcium levels were increased by 6 hours post-dose, achieved a maximum by 24 hours post-dose, and decreased gradually at 48 and 96 hours post-dose. Other clinical chemistry abnormalities, in group 3 and 4 animals included increased serum

proteins, cholesterol and kidney function parameters and decreased electrolytes and urine specific gravity. No toxicologically significant clinical chemistry abnormalities or notable increases in serum calcium were observed in group 2 animals.

[0126] There were no treatment-related changes observed in the ocular tissues on study days 22/23 and there were no treatment-related changes observed in the ECG and blood pressure data obtained on this study.

[0127] The most notable gross necropsy abnormalities occurred in group 4 animals that were found dead or were euthanized and included lesions in the digestive system and related organs; dark red omentum, reddened to dark red mucosa, red fluid in the small intestine and stomach, reddened to dark red mucosa in the esophagus and large intestine, stained and thickened gall bladder, a thrombus in the heart, dark red and mottled areas on the lungs, a reddened to dark red pancreas, a dark red thymus, thickened urinary bladder and a pale spleen. Gross abnormalities were less severe in group 3 animals; no notable gross abnormalities were observed in the group 2 animals.

[0128] The primary histopathological abnormality was dose related chronic interstitial nephritis: mild to moderate in group 3 animals and moderate to marked in group 4 animals. Other microscopic findings in these animals appeared to be secondary to chronic interstitial nephritis and included mineralization of various organs/tissues. No microscopic lesions were observed in the group 2 animals.

[0129] The highest values for serum calcium usually occurred within 24 hours post-dose and returned to baseline levels by the next pre-dose sampling interval. Selected data (males on Day 21) for serum calcium along with plasma calcitriol are shown in Figs. 5A-5C. These data show that the maximum plasma concentrations of calcitriol usually occurred well in advance of the maximum serum concentrations of calcium.

[0130] In summary, this study of semi-solid #3 administered orally to dogs once every 7 days to male and female Beagle dogs at 0, 1.0 and 5.0 (females) or 10.0 (males)  $\mu\text{g/kg}$  following the initial dose of 30.0  $\mu\text{g/kg}$  showed:



- The no observed adverse effect level was 0.1 µg/kg; the maximum tolerated dose was 1.0 µg/kg; mortality was seen at 30 µg/kg.
- Dose related lesions in the digestive system and related organs, reduced weight gain and decreased food consumption were seen in groups 3 and 4.
- Dose related chronic interstitial nephritis was seen in groups 3 and 4.

### EXAMPLE 13

#### Human Pharmacokinetic Study

- [0131] Pharmacokinetics of semi-solid #3 in humans was evaluated in a clinical trial. Patients received semi-solid #3 on this study at doses of calcitriol up to 90 µg. Preliminary pharmacokinetic results are discussed below.
- [0132] Blood samples were obtained pre-dose and at 0.5, 1.0, 1.5, 2, 3, 4, 6, 8, 12, 24, 48 and 72 hours post initial dose of semi-solid #3. Calcitriol levels were analyzed using a commercial radioimmunoassay, with limited validation for dilution integrity.
- [0133] Mean plasma concentration-time curves were plotted for each group (Fig. 6). Non-compartmental pharmacokinetic parameters were calculated for each subject and then averaged (Table 19). Baseline calcitriol values were subtracted from the post-dosing values to adjust for endogenous calcitriol.

Table 19: Semi-Solid #3 Pharmacokinetic Parameters by Dose Group

Dose, µg	C <sub>max</sub> , pg/mL (±SD)	T <sub>max</sub> , h (median and range)	AUC <sub>0-24h</sub> , pg·h/mL (±SD)	AUC <sub>0-48h</sub> , pg·h/mL (±SD)	AUC <sub>0-∞</sub> , pg·h/mL (±SD)	t <sub>1/2</sub> , h*
15.0 (n=3)	398.3 (12.9)	1.00 (1-1)	3665.7**	5627.3*** (637.1)	5464.8 (892.8)	8.9
30.0 (n=3)	898.8 (333.6)	1.50 (1.5-2)	6955.9 (2825.4)	9792.4 (2323.9)	11069.7*** (1406.4)	16.3***
60.0 (n=6)	2077.3 (533.3)	4.00 (1.5-4)	17480.6 (2989.7)	20999.4 (4762.5)	21795.0 (5124.8)	7.3
60.0 (n=4)	1918.4 (605.2)	1.3 (1-1.5)	17523.1 (1217.2)	20663.5 (1832.1)	24997.6 (4612.5)	8.6
75.0 (n=3)	1586.2 (328.6)	1.5 (1-4)	16499.1 (2343.8)	21159.1 (3406.0)	22690.4 (9209.4)	10.8
90.0 (n=3)	2858.7 (496.3)	1.5 (1-2)	23127.5 (5755.7)	28164.3 (8428.3)	29204.1 (9209.4)	8.8

\*harmonic mean, based on jackknife variance; \*\*n=1; \*\*\*n=2

[0134] Based on these data, pharmacokinetics of semi-solid #3 appear linear and predictable. There was no evidence of saturation of absorption.

#### EXAMPLE 14

##### Safety Results with Semi-Solid #3

[0135] The safety of semi-solid #3 in humans was evaluated in a clinical trial. As of May 8, 2002, 12 patients received semi-solid #3 on this study: 3 in group 1 (15 µg), 3 in group 2 (30 µg), and 6 in group 3 (60 µg). Preliminary pharmacokinetic results on the first 9 patients are discussed below.

[0136] No deaths have occurred. Thirty-four (34) adverse events occurred in 8 of the 9 patients; 20 of 34 adverse events were deemed possibly of probably related to semi-solid #3. One serious adverse event occurred in group 3 that was deemed not related by the Investigator. This patient developed a transient grade 1 fever on day 1 that prolonged hospitalization. Grade 2 or 3 adverse events deemed related to study drug are presented in Table 20.

Table 20: Grade 2 or 3 Adverse Events Deemed Related to Study Drug

Patient	Dose Group	Event	Severity	Comments
002-1002	60 µg	Hyperglycemia	Grade 2	-
		Hypoproteinemia	Grade 2	-
002-1003	60 µg	Constipation	Grade 2	-
		Hyponatremia	Grade 3	Sodium 127 meq/L on day 4; transient; no intervention

[0137] The preliminary results from the phase 1 trial with semi-solid #3 demonstrate:

The maximum tolerated dose of semi-solid #3 has not yet been determined in the phase 1 trial; additional patients are being evaluated in group 3 (60 µg).

Pharmacokinetics of semi-solid #3 appeared linear and predictable across the first three dose groups.

[0138] Having now fully described this invention, it will be understood by those of ordinary skill in the art that the same can be performed within a wide and equivalent range of conditions, formulations and other parameters without affecting the scope of the invention or any embodiment thereof. All patents, patent applications and publications cited herein are fully incorporated by reference herein in their entirety.

## WHAT IS CLAIMED IS:

1. A pharmaceutical composition comprising:

- (a) a lipophilic phase component,
- (b) one or more surfactants, and
- (c) an active vitamin D compound;

wherein said composition is an emulsion pre-concentrate, which, upon dilution with water in water to composition ratio of about 1:1 or more of said water, forms an emulsion having an absorbance of greater than 0.3 at 400 nm.

2. The pharmaceutical composition of claim 1, wherein said emulsion that is formed upon dilution of said emulsion pre-concentrate with water is a sub-micron droplet emulsion.

3. The pharmaceutical composition of claim 1, further comprising a hydrophilic phase component.

4. The pharmaceutical composition of claim 3, wherein said hydrophilic phase component comprises 1,2-propylene glycol.

5. The pharmaceutical composition of claim 4, wherein said hydrophilic phase component further comprises a C<sub>1-5</sub> alkanol.

6. The pharmaceutical composition of claim 5, wherein said C<sub>1-5</sub> alkanol is ethanol.

7. The pharmaceutical composition of claim 1, wherein said lipophilic phase component comprises a mono-, di- or triglyceride.

8. The pharmaceutical composition of claim 7, wherein said lipophilic phase component comprises a triglyceride.

9. The pharmaceutical composition of claim 8, wherein said triglyceride is selected from the group consisting of vegetable oils, fish oils, animal fats, hydrogenated vegetable oils, partially hydrogenated vegetable oils, synthetic triglycerides, modified triglycerides, fractionated triglycerides, and mixtures thereof.

10. The pharmaceutical composition of claim 1, wherein said one or more surfactants is a hydrophilic surfactant.

11. The pharmaceutical composition of claim 10, wherein said hydrophilic surfactant is a non-ionic hydrophilic surfactant.

12. The pharmaceutical composition of claim 11, wherein said non-ionic hydrophilic surfactant is selected from the group consisting of polyoxyethylene alkylethers; polyethylene glycol fatty acids esters; polyethylene glycol glycerol fatty acid esters; polyoxyethylene sorbitan fatty acid esters; polyoxyethylene-polyoxypropylene block copolymers; polyglycerol fatty acid esters; polyoxyethylene glycerides; polyoxyethylene vegetable oils; polyoxyethylene hydrogenated vegetable oils; reaction mixtures of polyols and at least one member of the group consisting of fatty acids, glycerides, vegetable oils, hydrogenated vegetable oils, and sterols; and mixtures thereof.

13. The pharmaceutical composition of claim 1, wherein said one or more surfactants is a lipophilic surfactant.

14. The pharmaceutical composition of claim 13, wherein said lipophilic surfactant is selected from the group consisting of alcohols; polyoxyethylene alkylethers; fatty acids; bile acids; glycerol fatty acid esters; acetylated glycerol fatty acid esters; lower alcohol fatty acids esters; polyethylene glycol fatty acids esters; polyethylene glycol glycerol fatty acid esters; polypropylene glycol fatty acid esters; polyoxyethylene glycerides; lactic acid

-53-

esters of mono/diglycerides; propylene glycol diglycerides; sorbitan fatty acid esters; polyoxyethylene sorbitan fatty acid esters; polyoxyethylene-polyoxypropylene block copolymers; trans-esterified vegetable oils; sterols; sugar esters; sugar ethers; sucroglycerides; polyoxyethylene vegetable oils; polyoxyethylene hydrogenated vegetable oils; reaction mixtures of polyols and at least one member of the group consisting of fatty acids, glycerides, vegetable oils, hydrogenated vegetable oils, and sterols; and mixtures thereof.

15. The pharmaceutical composition of claim 1, wherein said one or more surfactants is a surfactant containing a triglyceride.

16. The pharmaceutical composition of claim 1, wherein said surfactant is GELUCIRE.

17. The pharmaceutical composition of claim 1, comprising a single surfactant.

18. The pharmaceutical composition of claim 17, wherein said single surfactant is tocopherol PEG-1000 succinate (vitamin E TPGS).

19. The pharmaceutical composition of claim 1, further comprising at least one additive selected from the group consisting of an antioxidant, a bufferant, an antifoaming agent, a detackifier, a preservative, a chelating agent, a viscomodulator, a tonicifier, a flavorant, a colorant, an odorant, an opacifier, a suspending agent, a binder, a filler, a plasticizer, a thickening agent, and a lubricant.

20. The pharmaceutical composition of claim 19, wherein one of said additives is an antioxidant.

21. The pharmaceutical composition of claim 20, wherein said antioxidant is selected from the group consisting of butyl hydroxy anisole (BHA), and butyl hydroxy toluene (BHT).

22. The pharmaceutical composition of claim 1 adapted for oral administration.

23. The pharmaceutical composition of claim 22 in unit dosage form.

24. The pharmaceutical composition of claim 23 comprising 5-100  $\mu\text{g}$  of an active vitamin D compound per said unit dose.

25. The pharmaceutical composition of claim 24 comprising 10-75  $\mu\text{g}$  of an active vitamin D compound per said unit dose.

26. The pharmaceutical composition of claim 24, wherein said active vitamin D compound is calcitriol.

27. The pharmaceutical composition of claim 23, wherein said unit dosage form is a capsule.

28. The pharmaceutical composition of claim 27, wherein said capsule is a gelatin capsule.

29. The pharmaceutical composition of claim 27, wherein the total volume of ingredients present in said gelatin capsule is 10-1000  $\mu\text{L}$ .

30. The pharmaceutical composition of claim 1 in a form suitable for topical application.

31. The pharmaceutical composition of claim 1, wherein said lipophilic phase component is present in an amount of from 50 to 85% by weight based upon the total weight of the composition.

32. The pharmaceutical composition of claim 1, wherein said one or more surfactant is present in an amount of from 5 to 40% by weight based upon the total weight of the composition.

33. The pharmaceutical composition of claim 1, wherein said active vitamin D compound is present in an amount of from 0.01 to 15% by weight based upon the total weight of the composition.

34. The pharmaceutical composition of claim 3, wherein said hydrophilic phase component is present in an amount of from 5 to 15% by weight based upon the total weight of the composition.

35. The pharmaceutical composition of claim 1, wherein said composition is in a semisolid formulation.

36. The pharmaceutical composition of claim 35, wherein said lipophilic phase component is present in an amount of from 60 to 80% by weight based upon the total weight of the composition; said one or more surfactant is present in an amount of from 5 to 35% by weight based upon the total weight of the composition; and said active vitamin D compound is present in an amount of from 0.01 to 15% by weight based upon the total weight of the composition.

37. The pharmaceutical composition of claim 3, wherein said composition is in a liquid formulation.

38. The pharmaceutical composition of claim 37, wherein said lipophilic phase component is present in an amount of from 50 to 60% by weight



based upon the total weight of the composition, said one or more surfactant is present in an amount of from 4 to 25% by weight based upon the total weight of the composition, said active vitamin D compound is present in an amount of from 0.01 to 15% by weight based upon the total weight of the composition, and said hydrophilic phase component is present in an amount of from 5 to 10% by weight based upon the total weight of the composition.

39. A pharmaceutical emulsion composition comprising water and the pharmaceutical composition of claim 1.

40. The pharmaceutical composition of claim 1, wherein said active vitamin D compound is calcitriol.

41. A pharmaceutical sub-micron droplet emulsion composition comprising water and the pharmaceutical composition of claim 2.

42. An oral pharmaceutical composition comprising:

- (a) a lipophilic phase component,
- (b) one or more surfactants, and
- (c) an active vitamin D compound;

wherein said composition is an emulsion pre-concentrate, which, upon dilution with water in a water to composition ratio of about 1:1 or more of said water, forms an emulsion having an absorbance of greater than 0.3 at 400 nm.

43. The oral pharmaceutical composition of claim 42, wherein said emulsion that is formed upon dilution of said emulsion pre-concentrate with water is a sub-micron droplet emulsion.

44. The oral pharmaceutical composition of claim 42, further comprising a hydrophilic phase component.

-57-

45. A pharmaceutical composition comprising:

- (a) an active vitamin D compound,
- (b) a medium chain triglyceride,
- (c) tocopherol PEG-1000 succinate (vitamin E TPGS), and
- (d) GELUCIRE 44/14;

wherein said composition is an emulsion pre-concentrate, which, upon dilution with water in water to composition ratio of about 1:1 or more of said water, forms an emulsion having an absorbance of greater than 0.3 at 400 nm.

46. The pharmaceutical composition of claim 45, wherein said medium chain triglyceride is MIGLYOL 812.

47. The pharmaceutical composition of claim 46, wherein said active vitamin D compound is calcitriol.

48. The pharmaceutical composition of claim 47, wherein said calcitriol is present in an amount of from 0.01 to 0.05% by weight based upon the total weight of the composition, said MIGLYOL 812 is present in an amount of from 60 to 80% by weight based upon the total weight of the composition, said vitamin E TPGS is present in an amount of from 4 to 10% by weight based upon the total weight of the composition, and said GELUCIRE 44/14 is present in an amount of from 25 to 40% by weight based upon the total weight of the composition.

49. A method for the treatment or prevention of a hyperproliferative disease, said method comprising administering the pharmaceutical composition of claim 1 to a patient in need thereof.

50. The method of claim 49, wherein said hyperproliferative disease is cancer.

51. The method of claim 49, wherein the pharmaceutical composition is administered by pulse-dose, wherein said pulse-dose comprises the administration of said composition to a patient once every three to ten days.

52. A pharmaceutical composition comprising an active vitamin D compound, wherein said composition when administered to a subject provides a  $T_{\max}$  of less than about 6.0 hours and a  $T_{1/2}$  of less than about 25 hours.

53. The pharmaceutical composition of claim 52, wherein the  $T_{\max}$  is between about 1.0 hours and about 3.0 hours.

54. The pharmaceutical composition of claim 53, wherein the  $T_{\max}$  is between about 1.5 hours and about 2.0 hours.

55. The pharmaceutical composition of claim 52, wherein the  $T_{1/2}$  is between about 2 hours and about 10 hours.

56. The pharmaceutical composition of claim 55, wherein the  $T_{1/2}$  is between about 5 hours and about 9 hours.

57. The pharmaceutical composition of claim 52, wherein said composition when administered to a subject provides a  $C_{\max}$  of at least about 900 pg/mL.

58. The pharmaceutical composition of claim 57, wherein the  $C_{\max}$  is between about 900 and about 3000 pg/mL.

59. The pharmaceutical composition of claim 58, wherein the  $C_{\max}$  is between about 1500 and about 3000 pg/mL.

60. The pharmaceutical composition of claim 52, wherein said composition further comprises:

- i. a lipophilic phase component, and
- ii. one or more surfactants;

wherein said composition is an emulsion pre-concentrate, which, upon dilution with water in water to composition ratio of about 1:1 or more of said water, forms an emulsion having an absorbance of greater than 0.3 at 400 nm.

61. The pharmaceutical composition of claim 60, wherein said emulsion that is formed upon dilution of said emulsion pre-concentrate with water is a sub-micron droplet emulsion.

62. The pharmaceutical composition of claim 60, further comprising a hydrophilic phase component.

63. The pharmaceutical composition of claim 62, wherein said hydrophilic phase component comprises 1,2-propylene glycol.

64. The pharmaceutical composition of claim 63, wherein said hydrophilic phase component further comprises a C<sub>1-5</sub> alkanol.

65. The pharmaceutical composition of claim 64, wherein said C<sub>1-5</sub> alkanol is ethanol.

66. The pharmaceutical composition of claim 60, wherein said lipophilic phase component comprises a mono-, di- or triglyceride.

67. The pharmaceutical composition of claim 66, wherein said lipophilic phase component comprises a triglyceride.

68. The pharmaceutical composition of claim 67, wherein said triglyceride is selected from the group consisting of vegetable oils, fish oils, animal fats, hydrogenated vegetable oils, partially hydrogenated vegetable oils, synthetic triglycerides, modified triglycerides, fractionated triglycerides, and mixtures thereof.

69. The pharmaceutical composition of claim 60, wherein said one or more surfactants is a hydrophilic surfactant.

70. The pharmaceutical composition of claim 69, wherein said hydrophilic surfactant is a non-ionic hydrophilic surfactant.

71. The pharmaceutical composition of claim 70, wherein said non-ionic hydrophilic surfactant is selected from the group consisting of polyoxyethylene alkylethers; polyethylene glycol fatty acids esters; polyethylene glycol glycerol fatty acid esters; polyoxyethylene sorbitan fatty acid esters; polyoxyethylene-polyoxypropylene block copolymers; polyglycerol fatty acid esters; polyoxyethylene glycerides; polyoxyethylene vegetable oils; polyoxyethylene hydrogenated vegetable oils; reaction mixtures of polyols and at least one member of the group consisting of fatty acids, glycerides, vegetable oils, hydrogenated vegetable oils, and sterols; and mixtures thereof.

72. The pharmaceutical composition of claim 60, wherein said one or more surfactants is a lipophilic surfactant.

73. The pharmaceutical composition of claim 72, wherein said lipophilic surfactant is selected from the group consisting of alcohols; polyoxyethylene alkylethers; fatty acids; bile acids; glycerol fatty acid esters; acetylated glycerol fatty acid esters; lower alcohol fatty acids esters; polyethylene glycol fatty acids esters; polyethylene glycol glycerol fatty acid esters; polypropylene glycol fatty acid esters; polyoxyethylene glycerides; lactic acid

esters of mono/diglycerides; propylene glycol diglycerides; sorbitan fatty acid esters; polyoxyethylene sorbitan fatty acid esters; polyoxyethylene-polyoxypropylene block copolymers; trans-esterified vegetable oils; sterols; sugar esters; sugar ethers; sucroglycerides; polyoxyethylene vegetable oils; polyoxyethylene hydrogenated vegetable oils; reaction mixtures of polyols and at least one member of the group consisting of fatty acids, glycerides, vegetable oils, hydrogenated vegetable oils, and sterols; and mixtures thereof.

74. The pharmaceutical composition of claim 60, wherein said one or more surfactants is a surfactant containing a triglyceride.

75. The pharmaceutical composition of claim 60, wherein said surfactant is GELUCIRE.

76. The pharmaceutical composition of claim 60, comprising a single surfactant.

77. The pharmaceutical composition of claim 76, wherein said single surfactant is tocopherol PEG-1000 succinate (vitamin E TPGS).

78. The pharmaceutical composition of claim 60, further comprising at least one additive selected from the group consisting of an antioxidant, a bufferant, an antifoaming agent, a detackifier, a preservative, a chelating agent, a viscomodulator, a tonicifier, a flavorant, a colorant, an odorant, an opacifier, a suspending agent, a binder, a filler, a plasticizer, a thickening agent, and a lubricant.

79. The pharmaceutical composition of claim 78, wherein one of said additives is an antioxidant.

-62-

80. The pharmaceutical composition of claim 79, wherein said antioxidant is selected from the group consisting of butyl hydroxy anisole (BHA), and butyl hydroxy toluene (BHT).

81. The pharmaceutical composition of claim 60 adapted for oral administration.

82. The pharmaceutical composition of claim 81 in unit dosage form.

83. The pharmaceutical composition of claim 82 comprising 5-100  $\mu\text{g}$  of an active vitamin D compound per said unit dose.

84. The pharmaceutical composition of claim 83 comprising 10-75  $\mu\text{g}$  of an active vitamin D compound per said unit dose.

85. The pharmaceutical composition of claim 83, wherein said active vitamin D compound is calcitriol.

86. The pharmaceutical composition of claim 82, wherein said unit dosage form is a capsule.

87. The pharmaceutical composition of claim 86, wherein said capsule is a gelatin capsule.

88. The pharmaceutical composition of claim 87, wherein the total volume of ingredients present in said gelatin capsule is 10-1000  $\mu\text{L}$ .

89. The pharmaceutical composition of claim 60 in a form suitable for topical application.

90. The pharmaceutical composition of claim 60, wherein said lipophilic phase component is present in an amount of from 50 to 85% by weight based upon the total weight of the composition.

91. The pharmaceutical composition of claim 60, wherein said one or more surfactant is present in an amount of from 5 to 40% by weight based upon the total weight of the composition.

92. The pharmaceutical composition of claim 60, wherein said active vitamin D compound is present in an amount of from 0.01 to 15% by weight based upon the total weight of the composition.

93. The pharmaceutical composition of claim 62, wherein said hydrophilic phase component is present in an amount of from 5 to 15% by weight based upon the total weight of the composition.

94. The pharmaceutical composition of claim 60, wherein said composition is in a semisolid formulation.

95. The pharmaceutical composition of claim 94, wherein said lipophilic phase component is present in an amount of from 60 to 80% by weight based upon the total weight of the composition; said one or more surfactant is present in an amount of from 5 to 35% by weight based upon the total weight of the composition; and said active vitamin D compound is present in an amount of from 0.01 to 15% by weight based upon the total weight of the composition.

96. The pharmaceutical composition of claim 60, wherein said composition is in a liquid formulation.

97. The pharmaceutical composition of claim 96, wherein said lipophilic phase component is present in an amount of from 50 to 60% by weight



based upon the total weight of the composition, said one or more surfactant is present in an amount of from 4 to 25% by weight based upon the total weight of the composition, said active vitamin D compound is present in an amount of from 0.01 to 15% by weight based upon the total weight of the composition, and said hydrophilic phase component is present in an amount of from 5 to 10% by weight based upon the total weight of the composition.

98. A pharmaceutical emulsion composition comprising water and the pharmaceutical composition of claim 60.

99. The pharmaceutical composition of claim 52, wherein said active vitamin D compound is calcitriol.


100. A pharmaceutical sub-micron droplet emulsion composition comprising water and the pharmaceutical composition of claim 61.

101. A method for the treatment or prevention of a hyperproliferative disease, said method comprising administering the pharmaceutical composition of claim 52 to a patient in need thereof.


102. The method of claim 101, wherein said hyperproliferative disease is cancer.

103. The method of claim 101, wherein the pharmaceutical composition is administered by pulse-dose, wherein said pulse-dose comprises the administration of said composition to a patient once every three to ten days.


FIG. 1




2B




2A





3A




3B



**4A****4B**


5A

Calcitriol/Calcium Data for 0.1 ug/kg Males on Day 21




5B

Calcitriol/Calcium Data for 1 ug/kg Males on Day 21



5C

Calcitriol/Calcium Data for 10.0 ug/kg Males on Day 21






FIG. 6

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/38483

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/59

US CL : 514/167, 168

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/167, 168

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched :

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
EAST, USPATFUL, CAPLUS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5,486,509 A (JIMENEZ et al.) 23 January 1996 (23.01.1996), see the entire article, especially col. 9, example VIII.	1-103
X	US 5,502,224 A (EUGSTER et al.) 26 March 1996 (26.03.1996), see the entire article, especially col. 7, lines 14-49 and col. 16, line 58 - col. 17, line 45.	1-103

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

Special categories of cited documents:	
*A* document defining the general state of the art which is not considered to be of particular relevance	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*E* earlier application or patent published on or after the international filing date	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*O* document referring to an oral disclosure, use, exhibition or other means	*Z* document member of the same patent family
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

13 February 2003 (13.02.2003)

Date of mailing of the international search report

19 MAR 2003

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks

Box PCT

Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Valerie Bell-Hansford  
Barbara P. Badio, Ph.D.

Telephone No. 703-308-1235